



**B.A. PART- I
(SEMESTER - I)**

**MATHEMATICS : PAPER III
LINEAR ALGEBRA**

**Department of Distance Education
Punjabi University, Patiala**
(All Copyrights are Reserved)

Lesson No. :

SECTION - A

- 1.1 : RANK OF A MATRIX**
- 1.2 : ROW RANK, COLUMN RANK AND THEIR EQUIVALENCE**
- 1.3 : EIGEN VALUES AND EIGEN VECTORS**
- 1.4 : SYSTEM OF LINEAR EQUATIONS AND ITS CONSISTENCY**

SECTION - B

- 2.1 : VECTOR SPACES-I**
- 2.2 : VECTOR SPACES-II**
- 2.3 : LINEAR TRANSFORMATION-I**
- 2.4 : LINEAR TRANSFORMATION-II**

RANK OF A MATRIX

Objectives

- I. Introduction**
- II. Rank of a Matrix**
- III. Elementary Operations (or Transformations)**
- IV. Determination of Rank by Equivalent Matrix.**
- V. Computation of Inverse using Elementary**
- VI. Self Check Exercise**

I. Introduction

To understand the concept of rank of a matrix, firstly we take a look at the various types of matrices we have studied in our previous class.

(I) Transpose of a matrix : The matrix obtained from a given matrix A, by interchanging its rows and columns, is called the transpose of A and is generally denoted by A' or A^t or A^T .

Thus if $A = [a_{ij}]$ then (j, i) th element of A' is equal to (i, j) element of A

For example : If $A = \begin{bmatrix} 1 & -1 & 0 \\ 2 & 1 & 3 \end{bmatrix}$, then $A' = \begin{bmatrix} 1 & 2 \\ -1 & 1 \\ 0 & 3 \end{bmatrix}$

Remarks :

- (1) $(A')' = A$
- (2) $(A + B)' = A' + B'$
- (3) $(k A)' = k A'$, k being a complex number.
- (4) $(AB)' = B'A'$

(II) Symmetric and Skew-Symmetric Matrices

1. Any square matrix $A = [a_{ij}]$ is said to be a symmetric matrix if $a_{ij} = a_{ji}$ i.e., (i, j) th element of A is the same as the (j, i) th element of A. If we take the transpose of a symmetric matrix A, it is the same as A.

For example : $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}, \begin{bmatrix} 1 & -1 & 2 \\ -1 & 4 & 3 \\ 2 & 3 & 5 \end{bmatrix}$

2. Any square matrix $A = [a_{ij}]$ is said to be a skew-symmetric matrix if $a_{ij} = -a_{ji}$ i.e.

(i, j)th element is the same as the negative of the (j, i) element.

$\therefore$ for a skew-symmetric matrix A, $a_{ij} = -a_{ji}$

If we put $j = i$, we get $a_{ij} = -a_{ji}$ or $a_{ii} = 0$ i.e, every diagonal element of A is zero.

Examples of skew-symmetric matrix are

$$\begin{bmatrix} 0 & h & g \\ -h & 0 & f \\ -g & -f & 0 \end{bmatrix}, \begin{bmatrix} 0 & -5 \\ 5 & 0 \end{bmatrix}$$

(III) Conjugate and Tranjugate of a Matrix

The Matrix obtained by replacing the elements by A by its complex conjugates, is called the conjugate of A and is generally denoted by $\bar{A}$.

Thus, if $A = [a_{ij}]$, $\bar{A} = [\bar{a}_{ij}]$ where denotes the conjugate of a_{ij}

For example : if $A = \begin{bmatrix} 2+3i & 7-5i & 6+i \\ 5 & 2+3i & 1-2i \\ -3-5i & 0 & 2-5i \end{bmatrix}$,

$$\text{then } \bar{A} = \begin{bmatrix} 2-3i & 7+5i & 6-i \\ 5 & 2-3i & 1+2i \\ -3+5i & 0 & 2+5i \end{bmatrix}$$

If all the element of A are real, then $\bar{A} = A$.

Note $(\bar{\bar{A}}) = A$

Tranjugate of Matrix

The conjugate of the transpose of a matrix A is called tranjugate of A and is denoted by A^0 . Thus $A^0 = \overline{(A')}$

Clearly $\overline{(A')} = (\bar{A})'$

$$\therefore A^\theta = (\bar{A})'$$

For example : if

$$A = \begin{bmatrix} 2+3i & 6-i & 5+2i \\ 3 & 2 & -1+5i \\ 0 & 7-3i & -5+6i \end{bmatrix}$$

$$\text{then } A^\theta = \begin{bmatrix} 2-5i & 3 & 0 \\ 6+i & 2 & 7+3i \\ 5-2i & -1-5i & -5-6i \end{bmatrix}$$

Note : $(A^\theta)^\theta = A$.

(IV) Hermitian and Skew-Hermitian Matrices

(1) A square matrix $A = [a_{ij}]$ is said to be hermitian if $a_{ij} = \bar{a}_{ji}$ i.e., (i, j)th element is the conjugate of the (j, i)th element.

Now, $a_{ii} = \bar{a}_{ii} \therefore a_{ii} = \bar{a}_{ii}$ i.e., the conjugate of any diagonal element is the same element.

$\therefore$ every diagonal element must be real.

For example :

$$\begin{bmatrix} 2 & 5-6i & 3-4i \\ 5+6i & 0 & 1-2i \\ 3+4i & 1+2i & 7 \end{bmatrix}, \begin{bmatrix} 0 & a+ib & c+id \\ a-ib & 1 & m+in \\ c-id & m-in & p \end{bmatrix}$$

(2) A square matrix $A = [a_{ij}]$ is said to be Skew-hermitian if $a_{ij} = -\bar{a}_{ji}$ i.e., (i, j)th element is the negative conjugate of (j, i) element.

Again as $a_{ij} = -\bar{a}_{ji} \therefore a_{ii} = -\bar{a}_{ii}$ i.e., $a_{ii} + \bar{a}_{ii} = 0$.

$\therefore$ every diagonal element must be either zero or a purely imaginary number.

For example :

$$\begin{bmatrix} 4i & 4-3i & 6+5i \\ -4-3i & 0 & 2+7i \\ -6+5i & -2+7i & -9i \end{bmatrix}, \begin{bmatrix} 5i & 3-7i \\ -3-7i & 9i \end{bmatrix}$$

(v) Orthogonal Matrix

A square matrix P over the field of reals is said to be orthogonal if and only if $P'P = I$.

Now, if P is orthogonal, then $P'P = I = PP'$.

$$\Rightarrow |P'P| = |I| \Rightarrow |P'| |P| = I$$

$$\Rightarrow |P| |P| = I \Rightarrow |P|^2 = I$$

$$\Rightarrow |P| = \pm I \Rightarrow |P| \neq 0$$

$\Rightarrow P$ is invertible

$\therefore$ If P is orthogonal, then P is invertible.

$$\text{Also } P'P = I \Rightarrow P' = P^{-1}$$

$$\Rightarrow PP' = PP^{-1} \Rightarrow PP' = I$$

$\therefore P$ is orthogonal iff $P'P = PP' = I$ i.e, iff $P' = P^{-1}$.

For example :

$$\begin{bmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{bmatrix}$$

(vi) Unitary Matrix

A square matrix P over the field of complex numbers is said to be unitary if and only if $P^0P = I = PP^0$.

Now, if P is unitary, then $P^0P = I$

$$\Rightarrow |P^0P| = |I| \Rightarrow |P^0| |P| = 1 \Rightarrow |P| |P| = 1 \Rightarrow \therefore |P^0| = |P|$$

$$\Rightarrow |P|^2 = I \Rightarrow |P| \neq 0$$

$\Rightarrow P$ is invertible

$\therefore$ if P is unitary, then P is invertible.

$$\text{Also } P^0P = I \Rightarrow P^0 = P^{-1} \Rightarrow PP^0 = PP^{-1} \Rightarrow PP^0 = I$$

$$\therefore |P| = \pm 1$$

$\Rightarrow$ absolute value of a determinant of a unitary matrix is 1.

(vii) Similar Matrices

Let A and B be square matrices of order n over a field. Then A is said to be similar to B over F if and only if there exists an n-rowed invertible matrix C over F such that

$$AC = CB \text{ i.e. } B = C^{-1}AC \text{ or } A = CB^{-1}.$$

II. Rank of a Matrix

Definition : A number r is said to be rank of a non-zero matrix A if

- (i) there exists at least one minor of order r of A which does not vanish, and
- (ii) every minor of order (r + 1), if any, vanishes

The rank of a matrix A is denoted by $\rho(A)$.

$\therefore$ We have $\rho(A) = r$.

In other words, the rank of a non-zero matrix is the largest order of any non-vanishing minor of the matrix.

Remarks : (i) the rank of a zero matrix is zero i.e., $\rho(O) = 0$ where O is a zero matrix.

(ii) the rank of a non-singular matrix of order n is n,

(iii) $\rho(A) \leq r$, if every minor of order (r + 1) vanishes,

(iv) $\rho(A) \geq r$, if there is a minor of order r which does not vanish.

Some Important Results :-

Result 1: Prove that the rank of the transpose of a matrix A is the same as that of the original matrix A.

Proof . If $A = O$, then $A' = O$

$$\therefore \rho(A) = 0 \text{ and } \rho(A') = 0$$

$$\Rightarrow \rho(A') = \rho(A)$$

$\therefore$ result is true in the case in which A is a zero matrix.

Now we discuss the case when $A \neq O$.

Let r be the rank of the matrix $A = [a_{ij}]$ where A is of type $m \times n$.

$\therefore$ there exists at least one square submatrix R of order r such that $|R| \neq 0$.

Now R' is also a square submatrix of A' of order r.

$$\text{Also } |R'| = |R| \neq 0$$

$$\therefore \rho(A') \geq r$$

If possible, suppose $\rho(A') > r$

$$\text{We take } \rho(A') = r + 1 \Rightarrow \rho(A') \geq r + 1 \quad [\because (A')' = A]$$

$$\Rightarrow \rho(A) \geq r+1$$

$$[\because (A')' = A]$$

which is impossible as $\rho(A) = r$

$\therefore$ our supposition is wrong

$$\therefore \rho(A') \not\geq r$$

from (1), we have.

$$\rho(A') = r \Rightarrow \rho(A') = \rho(A).$$

Result 2 : Prove that $d\rho(\lambda A) = \rho(A)$ where λ is a non-zero scalar.

Proof : If $A = O$, then $\lambda A = O$

$$\therefore \rho(A) = 0 \text{ and } \rho(\lambda A) = 0$$

$$\therefore \rho(\lambda A) = \rho(A)$$

$\therefore$ result is true in the case in which A is a zero matrix.

Now we discuss the case when $A \neq O$.

Let r be the rank of the matrix $A = [a_{ij}]$ where A is of type $m \times n$.

$\therefore$ there exists at least one square submatrix R of order r such that $|R| \neq 0$

Now λR is a square submatrix of matrix λA of order r .

$$\therefore |\lambda R| = \lambda^r |R| \neq 0 \text{ as } \lambda \neq 0, |R| \neq 0$$

$$\therefore \rho(\lambda A) \geq r \text{ then } \rho(\lambda A) \geq r \quad \dots(1)$$

If possible, suppose $\rho(\lambda A) > r$

We take $\rho(\lambda A) > r+1$

$$\therefore \rho\left(\frac{1}{\lambda}(\lambda A)\right) \geq r+1 \quad [\because \text{of (1)}]$$

$$\Rightarrow \rho(A) \geq r+1, \text{ which is impossible as } \rho(A) = r$$

$\therefore$ Our supposition is wrong

$$\therefore \rho(\lambda A) = r \text{ or } \rho(\lambda A) = \rho(A)$$

Result 3 : If A is an n -rowed non-singular matrix, then prove that $d\rho(A^{-1}) = \rho(A)$.

Hence deduce that $\rho(\text{adj.}A) = \rho(A)$.

Proof : Here A is an n-rowed non-singular matrix

$$\therefore |A| \neq 0$$

$$\Rightarrow \rho(A) = n$$

$$\therefore A \text{ is non-singular}$$

$$\therefore A^{-1} \text{ exists and } AA^{-1} = I$$

$$\Rightarrow |AA^{-1}| = |I| \Rightarrow |A||A^{-1}| = 1 \Rightarrow |A^{-1}| \neq 0$$

$$\therefore A^{-1} \text{ is an n-rowed non-singular matrix}$$

$$\Rightarrow \rho(A^{-1}) = n \Rightarrow \rho(A^{-1}) = \rho(A)$$

Deduction

$$\rho(\text{adj.}A) = \rho\left(\frac{1}{|A|}\text{adj.}A\right) \quad \left[\because \rho(A) = \rho(\lambda A)\right]$$

$$= \rho(A^{-1}) = \rho(A) \Rightarrow \rho(\text{adj.}A) = \rho(A).$$

Problem 1 : Find the rank of the matrix $\begin{bmatrix} 1 & -1 & 3 & 6 \\ 1 & 3 & -3 & -4 \\ 5 & 3 & 3 & 11 \end{bmatrix}$.

Solution : Let $A = \begin{bmatrix} 1 & -1 & 3 & 6 \\ 1 & 3 & -3 & -4 \\ 5 & 3 & 3 & 11 \end{bmatrix}$

Since there does not exist any minor of order 4 or A

$$\therefore \rho(A) \leq 3 \quad \dots(1)$$

$$\text{Now } \begin{vmatrix} 1 & -1 & 6 \\ 1 & 3 & -4 \\ 5 & 3 & 11 \end{vmatrix} = 1 \begin{vmatrix} 3 & -4 \\ 3 & 11 \end{vmatrix} - (-1) \begin{vmatrix} 1 & -4 \\ 5 & 11 \end{vmatrix} + 6 \begin{vmatrix} 1 & 3 \\ 5 & 3 \end{vmatrix}$$

$\therefore$ there exists a minor of order 3 of A which does not vanish.

$$\therefore \rho(A) \geq 3$$

From (1) and (2), we get,

$$\rho(A) = 3.$$

III. Elementary Operations (or Transformations)

We can also determine the rank of a matrix by using some other methods which are based on the elementary transformations of a matrix, that includes :

- (1) The interchange of any two parallel lines.
- (2) The multiplication of all the elements of any line by any non-zero number.
- (3) The addition to the elements of any line, the corresponding elements of any other line multiplied by any number.

Note : An elementary transformation is called a row transformation or a column transformation according as it applies to rows or columns. Therefore, there are three row transformation and three column transformations.

Symbols used for the transformations

- (1) R_{ij} or $R_i \leftrightarrow R_j$ stands for the interchange of the i th and j th rows.
- (2) $R_i^{(c)}$ or $R_i \rightarrow cR_i$ stands for the multiplication of the i th row by $c \neq 0$.
- (3) $R_{ij}^{(k)}$ or $R_i \rightarrow R_i + kR_j$ stands for addition to the i th row, the product of the j th row by k . Similarly
- (4) C_{ij} or $C_i \leftrightarrow C_j$ stands for the interchange of i th and j th columns.
- (5) $C_i^{(c)}$ or $C_i \rightarrow cC_i$, stands for the multiplication of the elements of the i th column by $c \neq 0$.
- (6) $C_{ij}^{(k)}$ or $C_i \rightarrow C_i + kC_j$ stands for addition to the i th column, the product of the j th column by k .

Definition of Elementary Matrix

A matrix, obtained from a unit matrix, by subjecting it to a single elementary transformation is called an elementary matrix.

Remarks :

IV. Determination of Rank by Equivalent Matrix.

When an elementary transformation is applied to a matrix, it results into a matrix of the same order and same rank. The resulted matrix said to be equivalent to the given matrix and we use the symbol $\sim$ to mean.

Let A be any given matrix, Reduce the matrix to equivalent matrix by using the

following steps :

- (i) Use row or column transformations, if necessary, to obtain a non-zero element (preferably 1) in the first row and the first column of the given matrix.
- (ii) Divide the first row by this element, if it is not 1.
- (iii) Subtract suitable multiples of the first row from the other rows so as to obtain zeros in the remainder of the first column.
- (iv) Subtract suitable multiples of the first column from the other columns so as to get zeros in the remainder of the first row.
- (v) Repeat the steps (i) – (iv) starting with the elements in the second-row and the second column.
- (vi) Continue in this way down the "main diagonal" either until the end of the diagonal is reached or until all the remaining elements in the matrix are zero. The rank of this matrix, which is equivalent to the given matrix A, can be determined by inspection and consequently the rank of the given matrix A can be determined.

Problem 2 : Using elementary transformations, find the rank of the matrix

$$\begin{bmatrix} 1 & 3 & 2 \\ 4 & 6 & 5 \\ 3 & 5 & 4 \end{bmatrix}$$

Sol. Let $A = \begin{bmatrix} 1 & 3 & 2 \\ 4 & 6 & 5 \\ 3 & 5 & 4 \end{bmatrix}$

$$\sim \begin{bmatrix} 1 & 3 & 2 \\ 0 & -6 & -3 \\ 0 & -4 & -2 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - 4R_1, R_3 \rightarrow R_3 - 3R_1$$

$$\sim \begin{bmatrix} 1 & 3 & 2 \\ 0 & 2 & 1 \\ 0 & 2 & 1 \end{bmatrix}, \text{ by } R_2 \rightarrow -\frac{1}{3}R_2, R_3 \rightarrow -\frac{1}{2}R_3$$

$$\sim \begin{bmatrix} 1 & 3 & 2 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - R_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 0 & 0 & 0 \end{bmatrix}, \text{ by } C_2 \rightarrow C_2 - 3C_1, C_3 \rightarrow C_3 - 2C_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}, \text{ by } C_3 \rightarrow C_3 - \frac{1}{2} C_2$$

The rank of $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 0 \end{bmatrix}$ is 2 as minor $\begin{vmatrix} 1 & 0 \\ 0 & 2 \end{vmatrix}$ of order 2 does not vanish

$$\therefore \rho(A) = 2.$$

Problem 3 : Find the rank of the matrix

$$A = \begin{bmatrix} 1 & -1 & 1 & 5 \\ 2 & 1 & -1 & -2 \\ 3 & -1 & -1 & 7 \end{bmatrix}$$

Sol. $A = \begin{bmatrix} 1 & -1 & 1 & 5 \\ 2 & 1 & -1 & -2 \\ 3 & -1 & -1 & 7 \end{bmatrix}$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 5 \\ 0 & 3 & -3 & -12 \\ 0 & 2 & -4 & -8 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - 3R_1$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 5 \\ 0 & 1 & 1 & -4 \\ 0 & 2 & -4 & -8 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - R_3$$

$$\sim \begin{bmatrix} 1 & -1 & 1 & 5 \\ 0 & 1 & 1 & -4 \\ 0 & 0 & -6 & 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - 2R_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & -4 \\ 0 & 0 & -6 & 0 \end{bmatrix}, \text{ by } C_2 \rightarrow C_2 + C_1, C_3 \rightarrow C_3 - C_1, C_4 \rightarrow C_4 - 5C_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -6 & 0 \end{bmatrix}, \text{ by } C_3 \rightarrow C_3 - C_2, C_4 \rightarrow C_4 + 4C_2$$

The rank of $\begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & -6 & 0 \end{bmatrix}$ is 3 as the minor $\begin{vmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -6 \end{vmatrix} = -6 \neq 0$ of order 3 does not

vanish

$$\therefore \rho(A) = 3.$$

Note : Normal form of a Matrix : The normal form of matrix A can be

$$\begin{bmatrix} I_r & O \\ O & O \end{bmatrix}, [I_r \quad O], \begin{bmatrix} I_r \\ O \end{bmatrix}, \text{ where } I_r \text{ is identity matrix of order 'r'}.$$

Remarks : 1. Every non-zero matrix of rank r can, by a sequence of elementary

transformations, be reduced to the form $\begin{bmatrix} I_r & O \\ O & O \end{bmatrix}$ where I, is a r-rowed unit matrix.

2. Let A be any non-zero matrix of rank r. Then there exist non-singular matrices

$$P \text{ and } Q \text{ such that } PAQ = \begin{bmatrix} I_r & O \\ O & O \end{bmatrix}.$$

3. A non-singular matrix can be reduced to a unit matrix by a series of elementary transformations.

4. Every non-singular matrix is a product of elementary matrices.

5. The rank of a matrix is not altered by pre-multiplication or post-multiplication

of the matrix with any non-singular matrix.

6. The rank of a product of two matrices cannot exceed the rank of either matrix.

Problem 4 : Prove that the matrix $\begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 0 \\ 0 & 1 & 2 \end{bmatrix}$ is equivalent to I_3 .

Sol. Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 2 & 3 & 0 \\ 0 & 1 & 2 \end{bmatrix}$

$$\sim \begin{bmatrix} 1 & 2 & 3 \\ 0 & -1 & -6 \\ 0 & 1 & 2 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - 2R_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & -6 \\ 0 & 1 & 2 \end{bmatrix}, \text{ by } C_2 \rightarrow C_2 - 2C_1, C_3 \rightarrow C_3 - 3C_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 6 \\ 0 & 1 & 2 \end{bmatrix}, \text{ by } R_2 \rightarrow -R_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 6 \\ 0 & 0 & -4 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - R_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -4 \end{bmatrix}, \text{ by } C_3 \rightarrow C_3 - 6C_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ by } R_3 \rightarrow -\frac{1}{4} R_3$$

$$\therefore A \sim I_3$$

$\therefore$ given matrix is equivalent to I_3 .

Problem 5 : If $A = \begin{bmatrix} 1 & 1 & 1 \\ 3 & 1 & 1 \\ 1 & -1 & -1 \end{bmatrix}$, then find the matrices P and Q such that PAQ is in

the normal form. Hence find the rank of the matrix A.

Sol. Here $A = \begin{bmatrix} 1 & 1 & 1 \\ 3 & 1 & 1 \\ 1 & -1 & -1 \end{bmatrix}$,

We have $A = I A I$

$$\therefore \begin{bmatrix} 1 & 1 & 1 \\ 3 & 1 & 1 \\ 1 & -1 & -1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & -1 \\ 3 & 1 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ by } R_2 \leftrightarrow R_3$$

$$\Rightarrow \begin{bmatrix} 1 & 1 & 1 \\ 0 & -2 & -2 \\ 0 & -2 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 0 & 1 \\ -3 & 1 & 0 \end{bmatrix} A \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

by $R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - 3R_1$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & -2 & -2 \\ 0 & -2 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ -1 & 0 & 1 \\ -3 & 1 & 0 \end{bmatrix} A \begin{bmatrix} 1 & -1 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix},$$

by $C_2 \rightarrow C_2 - C_1, C_3 \rightarrow C_3 - C_1$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & -2 & -2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 0 & -\frac{1}{2} \\ -3 & 1 & 0 \end{bmatrix} A \begin{bmatrix} 1 & -1 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ by } R_2 \rightarrow -\frac{1}{2}R_2$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 1 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 0 & -\frac{1}{2} \\ -2 & 1 & -1 \end{bmatrix} A \begin{bmatrix} 1 & -1 & -1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \text{ by } R_3 - R_3 + 2R_2$$

$$\Rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 0 & -\frac{1}{2} \\ -2 & 1 & -1 \end{bmatrix} A \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}, \text{ by } C_3 \rightarrow C_3 - C_2$$

$$\Rightarrow \begin{bmatrix} I_2 & O \\ O & O \end{bmatrix} = PAQ$$

$$\text{where } P = \begin{bmatrix} 1 & 0 & 0 \\ \frac{1}{2} & 0 & -\frac{1}{2} \\ -2 & 1 & -1 \end{bmatrix}, Q = \begin{bmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \end{bmatrix}$$

$\therefore$ PAQ is in the normal form and $\rho(A) = 2$.

Problem 6 : Reduce the matrix $\begin{bmatrix} 2 & -1 & 0 & 4 \\ 1 & 3 & 5 & -3 \\ 3 & -5 & -5 & 11 \\ 6 & 4 & 10 & 2 \end{bmatrix}$ to normal form. Hence find the

rank of the matrix.

Solution : $A = \begin{bmatrix} 2 & -1 & 0 & 4 \\ 1 & 3 & 5 & -3 \\ 3 & -5 & -5 & 11 \\ 6 & 4 & 10 & 2 \end{bmatrix}$

$$\sim \begin{bmatrix} 1 & 3 & 5 & -3 \\ 2 & -1 & 0 & 4 \\ 3 & -5 & -5 & 11 \\ 6 & 4 & 10 & 2 \end{bmatrix}, \text{by } R_1 \leftrightarrow R_2$$

$$\sim \begin{bmatrix} 1 & 3 & 5 & -3 \\ 0 & -7 & 10 & 10 \\ 0 & -14 & -20 & 20 \\ 0 & -14 & -20 & 20 \end{bmatrix}, \text{by } R_1 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - 3R_1, R_4 \rightarrow R_4 - 6R_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & -7 & 10 & 10 \\ 0 & -14 & -20 & 20 \\ 0 & -14 & -20 & 20 \end{bmatrix}, \text{by } C_2 \rightarrow C_2 - 3C_1, C_3 \rightarrow C_3 - 5C_1, C_4 \rightarrow C_4 + 3C_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 2 & 2 & 2 \\ 0 & 2 & 2 & 2 \end{bmatrix}, \text{by } C_2 \rightarrow -\frac{1}{7}C_2, C_3 \rightarrow -\frac{1}{10}C_3, C_4 \rightarrow \frac{1}{10}C_4$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{by } R_3 \rightarrow R_3 - 2R_2, R_4 \rightarrow R_4 - 2R_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{ by } C_3 \rightarrow C_3 - C_2, C_4 \rightarrow C_4 - C_2$$

$$\sim \begin{bmatrix} I_2 & O \\ O & O \end{bmatrix}$$

$$\therefore \text{rank}(A) = 2$$

V. Computation of Inverse using Elementary Transformations

We can understand this computation of finding inverse with the help of following example:

If we are to find inverse of A, we write $A = I A$ and go on performing row transformations on the product and the prefactor of A till we reach the result $I = BA$, then B is the inverse of A.

Problem 7 : Using elementary operations, find inverse of the matrix:

$$A = \begin{bmatrix} 2 & 3 & 1 \\ -3 & 5 & 1 \\ 1 & 7 & 2 \end{bmatrix}$$

Solution : $A = \begin{bmatrix} 2 & 3 & 1 \\ -3 & 5 & 1 \\ 1 & 7 & 2 \end{bmatrix}$

Now $A = I A$

$$\therefore \begin{bmatrix} 2 & 3 & 1 \\ -3 & 5 & 1 \\ 1 & 7 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} A$$

$$\therefore \begin{bmatrix} 1 & 7 & 2 \\ -3 & 5 & 1 \\ 2 & 3 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{bmatrix} A, \text{ by } R_2 \leftrightarrow R_3$$

$$\therefore \begin{bmatrix} 1 & 7 & 2 \\ 0 & 26 & 7 \\ 0 & -11 & -3 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 3 \\ 1 & 0 & -2 \end{bmatrix} \text{A, by } R_2 \rightarrow R_2 + 3R_1, R_3 \rightarrow R_3 - 2R_1$$

$$\therefore \begin{bmatrix} 1 & 7 & 2 \\ 0 & 26 & 7 \\ 0 & -286 & -78 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 3 \\ 26 & 0 & -52 \end{bmatrix} \text{A, by } R_3 \rightarrow 26R_3$$

$$\therefore \begin{bmatrix} 1 & 7 & 2 \\ 0 & 26 & 7 \\ 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 3 \\ 26 & 11 & -19 \end{bmatrix} \text{A, by } R_3 \rightarrow R_3 + 11R_2$$

$$\therefore \begin{bmatrix} 1 & 7 & 0 \\ 0 & 26 & 0 \\ 0 & 0 & -1 \end{bmatrix} = \begin{bmatrix} 52 & 22 & -37 \\ 182 & 78 & -130 \\ 26 & 11 & -19 \end{bmatrix} \text{A, by } R_1 \rightarrow R_1 + 2R_3, R_2 \rightarrow R_2 + 7R_3$$

$$\therefore \begin{bmatrix} 1 & 7 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 52 & 22 & -37 \\ 7 & 3 & -5 \\ -26 & -11 & 19 \end{bmatrix} \text{A, by } R_2 \rightarrow \frac{1}{26}R_2, R_3 \rightarrow -R_3$$

$$\therefore \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & 1 & -2 \\ 7 & 3 & -5 \\ -26 & -11 & 19 \end{bmatrix} \text{A, by } R_1 \rightarrow R_1 - 7R_2$$

$$\therefore I = A^{-1} A$$

$$\therefore A^{-1} = \begin{bmatrix} 3 & 1 & -2 \\ 7 & 3 & -5 \\ -26 & -11 & 19 \end{bmatrix}.$$

VI. Self Check Exercise

1. Show that the matrix $A = \begin{bmatrix} a + ic & -b + id \\ b + id & a - ic \end{bmatrix}$ is unitary if and only if

$$a^2 + b^2 + c^2 + d^2 = 1.$$

- 2.** (i) If P, Q are unitary, prove that QP is also unitary.
(ii) If P, Q are orthogonal, prove that QP is also orthogonal.
- 3.** If A is an orthogonal matrix, then A' and A^{-1} are also orthogonal.

4. Find the rank of the matrix $\begin{bmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$.

5. Find the rank of each matrix $\begin{bmatrix} 0 & 6 & 6 & 1 \\ -8 & 7 & 2 & 3 \\ -2 & 3 & 0 & 1 \\ -3 & 2 & 1 & 1 \end{bmatrix}$.

6. Find the rank of the matrix $\begin{bmatrix} 1 & 2 & -3 & -1 \\ 3 & -4 & 1 & 2 \\ 5 & 2 & 1 & 3 \end{bmatrix}$.

7. Find the rank of the matrix $\begin{bmatrix} 2 & 3 & 4 \\ 3 & 1 & 2 \\ -1 & 2 & 2 \end{bmatrix}$, using equivalent matrix.

8. Find the rank of the matrix $\begin{bmatrix} 3 & 4 & 1 & 2 \\ 3 & 2 & 1 & 4 \\ 7 & 6 & 2 & 5 \end{bmatrix}$, using equivalent matrix.

9. For the matrix $A = \begin{bmatrix} 1 & 1 & 2 \\ 1 & 2 & 3 \\ 0 & -1 & -1 \end{bmatrix}$, find two non-singular matrices P and

Q such that PAQ is in the normal form and hence find out rank of matrix A.

10. Reduce the matrix $\begin{bmatrix} 3 & -2 & 1 \\ 2 & -1 & 3 \\ 1 & -2 & 1 \end{bmatrix}$ to the form I_3 and find rank.

11. Use elementary transformation to find the inverse of

$$(i) \begin{bmatrix} 1 & 1 & 1 \\ 1 & -1 & 1 \\ 2 & 1 & -1 \end{bmatrix} \quad (ii) \begin{bmatrix} 1 & 3 & 2 \\ 0 & 4 & 1 \\ 5 & 2 & 3 \end{bmatrix}$$

ROW RANK, COLUMN RANK AND THEIR EQUIVALENCE

Objectives

- I. Row and Column Rank of a Matrix.**
- II. Linear Dependence|Independence of Vectors**
- III. Equality of Row Rank and column Rank Row Rank and column Rank:**
- IV. Self check Exercise**

I. Row and Column Rank of a Matrix.

Firstly, we define the echelon form of a matrix:

A matrix A is said to be a row (column) equivalent to a matrix B if B can be obtained from A after a finite number of elementary row (column) operations, and we write $A^R B$ or $A^C B$.

Definition (Echelon Form):

A matrix $A = [a_{ij}]$ is said to be in the echelon form if

- (i) The zero rows (columns) of A occur below all the non-zero rows (columns) of A
- (ii) The number of zeros before the first non-zero element in a row (column) is less than the number of such zeros in the next row (column).
- (iii) If $R_1, R_2, \dots$ are non-zero rows (columns) of A, then first non-zero entry in these rows (columns) is 1. The Moreover Matrix is in (column) row reduced echelon form in addition to the above conditions, if a column (row) contains the first non-zero entry of any row(column), then every other entry in that column (row) is zero.

Row and column Rank of a Matrix

Let A be any matrix. Then Row rank of A, denoted by $\rho_R(A)$, is defined as the number of non-zero rows in a row echelon form of A.

Similarly, column rank of A, denoted by $\rho_C(A)$, is defined as the number of non-zero column in a column echelon form of A.

Problem 1 : Find the row rank and column rank of $\begin{bmatrix} 1 & 3 & 2 & 4 \\ 5 & 2 & 0 & 1 \\ 3 & -4 & -4 & -7 \\ -7 & 5 & 6 & 10 \end{bmatrix}$.

Solution : Let $A = \begin{bmatrix} 1 & 3 & 2 & 4 \\ 5 & 2 & 0 & 1 \\ 3 & -4 & -4 & -7 \\ -7 & 5 & 6 & 10 \end{bmatrix}$

$$\sim \begin{bmatrix} 1 & 3 & 2 & 4 \\ 0 & -13 & -10 & -19 \\ 0 & -13 & -10 & -19 \\ -0 & 26 & 20 & 38 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - 5R_1, R_3 \rightarrow R_3 - 3R_1, R_4 \rightarrow R_4 + 7R_1$$

$$\sim \begin{bmatrix} 1 & 3 & 2 & 4 \\ 0 & -13 & -10 & -19 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - R_2, R_4 \rightarrow R_4 + 2R_2$$

$$\sim \begin{bmatrix} 1 & 3 & 2 & 4 \\ 0 & 1 & \frac{10}{13} & \frac{19}{13} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}, \text{ by } R_2 \rightarrow -\frac{1}{13}R_2$$

Which is in row-echelon form.

Since there are two non-zero rows in the row-echelon form

$\therefore$ row rank of A is 2

$$\text{Now } A = \begin{bmatrix} 1 & 3 & 2 & 4 \\ 5 & 2 & 0 & 1 \\ 3 & -4 & -4 & -7 \\ -7 & 5 & 6 & 10 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 5 & -13 & -10 & -19 \\ 3 & -13 & -10 & -19 \\ -7 & 26 & 20 & 38 \end{bmatrix}, \text{by } C_2 \rightarrow C_2 - 3C_1, C_3 \rightarrow C_3 - 2C_1, C_4 \rightarrow C_4 - 4C_1$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 5 & 1 & -10 & -19 \\ 3 & 1 & -10 & -19 \\ -7 & -2 & 20 & 38 \end{bmatrix}, \text{by } C_2 \rightarrow C_2 - \frac{1}{13}C_2$$

$$\sim \begin{bmatrix} 1 & 0 & 0 & 0 \\ 5 & 1 & 0 & 0 \\ 3 & 1 & 0 & 0 \\ -7 & -2 & 0 & 0 \end{bmatrix}, \text{by } C_3 \rightarrow C_3 + 10C_2, C_4 \rightarrow C_4 + 15C_2$$

which is column echelon form having two non-zero column.

$\therefore$ column rank = 2.

In order to understand the concept of row rank and column rank more deeply, we must have the knowledge of law vectors and column vectors.

II. Linear Dependence | Independence of Vectors

Definition (n-vector) : An ordered tuple of n numbers is called n-vector.

For example : $\{x_1, x_2, \dots, x_n\}$ is an n-vector.

Linear Dependence of Vectors : A set $V_1, V_2, \dots, V_t$ of vectors is said to be linearly dependent set, if there exists t scalars $p_1, p_2, \dots, p_t$, not all zero, such that $p_1V_1 + p_2V_2 + \dots + p_tV_t = O$, where O is a n-vector with all components zero.

Any set of vectors, which is not linearly dependent, is called linearly independent.

i.e. a set $V_1, V_2, \dots, V_t$ of n-vectors is said to be linearly independent if every

relation of the form $p_1V_1 + p_2V_2 + \dots + p_tV_t = O$ implies $p_1 = p_2 = \dots = p_t = 0$.

Linear Combination of Vectors :

A vector V is said to be a linear combination of the vectors

$V_1, V_2, \dots, V_t$ if $V = p_1V_1 + p_2V_2 + \dots + p_tV_t$, where $p_1, p_2, \dots, p_t$ are scalars.

Result 1 : If a set of vectors is linearly dependent, show that at least one member of the set is a linear combination of the remaining members.

Proof : Let $V_1, V_2, \dots, V_t$ be any linearly dependent set.

$$\therefore \text{the relation } p_1V_1 + p_2V_2 + \dots + p_tV_t = O$$

implies that at least one of $p_1, p_2, \dots, p_t$ is non-zero

Let p_1 be non-zero

$$\text{Now } p_1V_1 = -p_2V_2 - p_3V_3 - \dots - p_tV_t$$

$$\therefore V_1 = \left(-\frac{p_2}{p_1}\right)V_2 + \left(-\frac{p_3}{p_1}\right)V_3 + \dots + \left(-\frac{p_t}{p_1}\right)V_t$$

The relation (1) shows that V_1 is a linear combination of $V_2, V_3, \dots, V_t$.

Result 2 : If η is a linear combination of the set $\{V_1, V_2, \dots, V_r\}$, then the set

$\{\eta, V_1, V_2, \dots, V_r\}$ is linearly dependent.

Proof : Since η is a linear combination of $V_1, V_2, \dots, V_r$

$$\therefore \eta = k_1V_1 + k_2V_2 + \dots + k_rV_r$$

$$\Rightarrow \eta - k_1V_1 - k_2V_2 - \dots - k_rV_r = O$$

Now at least one of the coefficients i.e. of η is non-zero.

$$\therefore \text{set } \eta, V_1, V_2, \dots, V_r \text{ is L.D.}$$

Result 3 : Prove that every super set of a linearly dependent set is linearly dependent.

Proof : Let $\{V_1, V_2, \dots, V_p, V_{p+1}, \dots, V_r\}$ be a super set of a linearly dependent set

$\{V_1, V_2, \dots, V_p\}$. Since $\{V_1, V_2, \dots, V_p\}$ is linearly dependent set

$\therefore$ there exist scalars $k_1, k_2, \dots, k_p$ (not all zero) such that

$$k_1V_1 + k_2V_2 + \dots + k_pV_p = O$$

It can be re-written as

$k_1V_1 + k_2V_2 + \dots + k_pV_p + \dots + k_rV_r = O$, where $k_1, k_2, \dots, k_p, \dots, k_r$ and not all zero.

$\therefore$ set $\{V_1, V_2, \dots, V_p, \dots, V_r\}$ is L.D.

Brief Outline of Vector Space :

In this lesson we briefly explain what a vector space is? The detailed study of vector space will be done in lesson 0.5.

Definition : The n-vector Space : The set of all n-vectors over a field F, to be denoted by $V_n(F)$, is called the n-vector space over F.

Sub-space of n-vector Space V_n

Any non-zero empty set, S of vectors of $V_n(F)$ is called a subspace of $V_n(F)$, if when

- (i) V_1, V_2 are any two members of S, then $V_1 + V_2$ is also a member of S and
- (ii) If V is a member of S and k is a member of F, then kV is also a member of S.

Subspace Spanned by a Set of Vectors

Let $V_1, V_2, \dots, V_t$ be a set of n-vectors.

The set of all linear combinations of the above set is called a subspace spanned by the set of vectors $V_1, V_2, \dots, V_t$.

Basis of a Subspace

A set of vectors is said to be the basis of a subspace, if

- (i) the subspace is spanned by the set and
- (ii) the set is linearly independent.

Dimension of a subspace

The number of vectors in any basis of a subspace is called the dimension of the subspace.

Another Method to check for the Linear Dependence of Vectors :

Let $V_1 = (b_{11}, b_{12}, \dots, b_{1n}), V_2 = (b_{21}, b_{22}, \dots, b_{2n})$

.....
.....

$V_n = (b_{n1}, b_{n2}, \dots, b_{nn})$ be a vectors of the vector space.

By definition these vectors are L.D. vectors iff there exists scalars $\alpha_1, \alpha_2, \dots, \alpha_n \in F$,

not all zero such that $\alpha_1V_1 + \alpha_2V_2 + \dots + \alpha_nV_n = O$

$$\Rightarrow \alpha_1(b_{11}, b_{12}, \dots, b_{1n}) + \alpha_2(b_{21}, b_{22}, \dots, b_{2n}) + \dots + \alpha_n(b_{n1}, \dots, b_{nn}) = O$$

$$\Rightarrow (\alpha_1b_{11} + \alpha_2b_{21} + \dots + \alpha_nb_{n1}, \alpha_1b_{12} + \alpha_2b_{22} + \dots$$

$$\begin{aligned} \therefore \quad & \alpha_1 \mathbf{b}_{11} + \alpha_2 \mathbf{b}_{21} + \dots + \alpha_n \mathbf{b}_{n1} = 0 \\ & \alpha_1 \mathbf{b}_{12} + \alpha_2 \mathbf{b}_{22} + \dots + \alpha_n \mathbf{b}_{n2} = 0 \\ & \dots\dots\dots \\ & \alpha_1 \mathbf{b}_{1n} + \alpha_2 \mathbf{b}_{2n} + \dots + \alpha_n \mathbf{b}_{nn} = 0 \end{aligned}$$

$$\text{i.e.,} \quad \text{iff} \quad \begin{vmatrix} b_{11} & b_{21} & \dots & b_{n1} \\ b_{12} & b_{22} & \dots & b_{n2} \\ \dots & \dots & \dots & \dots \\ b_{1n} & b_{2n} & \dots & b_{nn} \end{vmatrix} = 0$$
$$\text{iff } \begin{vmatrix} b_{11} & b_{21} & \dots & b_{n1} \\ b_{12} & b_{22} & \dots & b_{n2} \\ \dots & \dots & \dots & \dots \\ b_{1n} & b_{2n} & \dots & b_{nn} \end{vmatrix} = 0$$

$$\text{or} \quad \text{iff} \quad \begin{vmatrix} b_{11} & b_{21} & \dots & b_{1n} \\ b_{21} & b_{22} & \dots & b_{2n} \\ \dots & \dots & \dots & \dots \\ b_{n1} & b_{n2} & \dots & b_{nn} \end{vmatrix} = 0$$

$\Rightarrow \exists$ scalars α_1, α_2 and α_3 such that

$$\begin{aligned}
(1, -3, 5) &= \alpha_1 (1, 2, 1) + \alpha_2 (1, 1, -1) + \alpha_3 (4, 5, -2) \\
&= (\alpha_1, 2\alpha_1, \alpha_1) + (\alpha_2, \alpha_2 - \alpha_2) + (4\alpha_3, 5\alpha_3, -2\alpha_3) \\
&= (\alpha_1 + \alpha_2 + 4\alpha_3, 2\alpha_1 + \alpha_2 + 5\alpha_3, \alpha_1 - \alpha_2 - 2\alpha_3)
\end{aligned}$$

By equality of vectors, we must have

$$\alpha_1 + \alpha_2 + 4\alpha_3 = 1$$

$$2\alpha_1 + \alpha_2 + 5\alpha_3 = -3$$

$$\alpha_1 - \alpha_2 - 2\alpha_3 = 5$$

Adding (1) and (3), we get,

$$2\alpha_1 + 2\alpha_3 = 6$$

$$\Rightarrow \alpha_1 + \alpha_3 = 3$$

and adding (2) and (3), we get,

$$3\alpha_1 + 3\alpha_3 = 2$$

$$\Rightarrow \alpha_1 + \alpha_3 = \frac{2}{3}$$

From (4) and (5), it is clear that we cannot find α_1 and α_2 and so α_3 .

$\therefore$ our supposition is wrong.

Hence $(1, -3, 5)$ does not belong to the Linear Space of S.

Problem 3 : Is the system of vectors $[-1, 1, 2]$, $[2, -3, 1]$, $[10, -1, 0]$ linearly dependent ?

Sol. Given vectors are

$$V_1 = [-1, 1, 2], V_2 = [2, -3, 1], V_3 = [10, -1, 0]$$

Consider the relation

$$k_1 V_1 + k_2 V_2 + k_3 V_3 = O$$

$$\text{or } k_1 [-1, 1, 2] + k_2 [2, -3, 1] + k_3 [10, -1, 0] = [0, 0, 0]$$

$$\therefore -k_1 + 2k_2 + 10k_3 = 0 \quad \dots (1)$$

$$k_1 - 3k_2 - k_3 = 0 \quad \dots (2)$$

$$2k_1 + k_2 = 0 \quad \dots (3)$$

From (3), $k_2 = -2k_1$

$$\begin{aligned}
\therefore \quad & \text{from (2), } k_1 + 6k_1 - k_3 = 0 \Rightarrow k_3 = 7k_1 \\
\therefore \quad & \text{from (1), } -k_1 - 4k_1 + 70k_1 = 0 \Rightarrow 65k_1 = 0 \Rightarrow k_1 = 0 \\
\therefore \quad & k_2 = 0, k_3 = 0 \\
\therefore \quad & k_1 = k_2 = k_3 = 0 \\
\therefore \quad & k_1V_1 + k_2V_2 + k_3V_3 = O \Rightarrow k_1 = k_2 = k_3 = 0 \\
\therefore \quad & \text{given set of vectors is L.I.}
\end{aligned}$$

Problem 4 : Find the value of k so that the vectors

$$\begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ -2 \end{bmatrix} \text{ and } \begin{bmatrix} k \\ 0 \\ 1 \end{bmatrix} \text{ are L.D.}$$

Sol. Let a, b, c be scalars, not all zero, such that

$$a \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} + b \begin{bmatrix} 1 \\ 2 \\ -2 \end{bmatrix} + c \begin{bmatrix} k \\ 0 \\ 1 \end{bmatrix} = O$$

where O is 3×1 zero matrix

$$\Rightarrow \begin{bmatrix} a \\ -a \\ 3a \end{bmatrix} + \begin{bmatrix} b \\ 2b \\ -2b \end{bmatrix} + \begin{bmatrix} ck \\ 0 \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a + b + ck \\ -a + 2b \\ 3a - 2b + c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore \quad a + b + ck = 0$$

$$-a + 2b = 0$$

$$3a - 2b + c = 0$$

From (2), we get $a = 2b$

$$\therefore 3 \Rightarrow 3a - a + c = 0 \Rightarrow c = -2a$$

Put the values of c and b in (1), we have

$$a + \frac{a}{2} - 2ak = 0 \Rightarrow \frac{3a}{2} - 2ak = 0 \Rightarrow 2a\left(\frac{3}{4} - k\right) = 0$$

But $a \neq 0$

[as if $a = 0$ then $b = 0$ and $c = 0$ which implies the given vectors are L.I.]

$$\Rightarrow \frac{3}{4} - k = 0 \Rightarrow k = \frac{3}{4}.$$

III. Equality of Row Rank and column Rank Row Rank and column Rank:

If A is any $m \times n$ matrix, then

(i) the space spanned by the set of m rows is called row space of A and the number of independent row vectors is called the row rank of A .

(ii) the space spanned by the set of n columns is called Column Space of A and the number of independent column vectors is called the column rank of A .

In other words, Column rank of any matrix A is the maximum number of linearly independent columns of A .

Result 4 : Prove that pre-multiplication by a non-singular matrix does not alter the row rank of a matrix.

Proof : Let $A = \begin{bmatrix} R_1 \\ R_2 \\ \vdots \\ R_m \end{bmatrix}$ be $m \times n$ matrix and

$$P = \begin{bmatrix} p_{11} & p_{12} & \dots & p_{1m} \\ p_{21} & p_{22} & \dots & p_{2m} \\ \dots & \dots & \dots & \dots \\ p_{m1} & p_{m2} & \dots & p_{mm} \end{bmatrix} \text{ be } m \times n \text{ non-singular matrix}$$

$$\text{Let } B = PA = \begin{bmatrix} p_{11} & p_{12} & \dots & p_{1m} \\ p_{21} & p_{22} & \dots & p_{2m} \\ \dots & \dots & \dots & \dots \\ p_{m1} & p_{m2} & \dots & p_{mm} \end{bmatrix} \begin{bmatrix} R_1 \\ R_2 \\ \vdots \\ R_m \end{bmatrix}$$

Note : Similarly we can prove that $AS = \begin{bmatrix} L & O \end{bmatrix}$, where L is an $m \times s$ matrix, s being the column rank of A .

Result 6 : Prove that the row rank of a matrix is the same as its rank.

Proof : Let r be the rank and s be the row rank of $m \times n$ matrix A .

Since s is row rank of A

$\therefore$ there exists a non-singular matrix R such that

$$RA = \begin{bmatrix} K \\ O \end{bmatrix}, \text{ where } K \text{ is } s \times n \text{ matrix}$$

since each minor of order $(s + 1)$ of the matrix RA involves at least are row of zeros

$$\therefore \rho(RA) \leq s$$

$$\therefore r \leq s \quad \dots(1)$$

Since r is rank of A

$\therefore$ there exists a non-singular matrix P such that

$$PA = \begin{bmatrix} G \\ O \end{bmatrix}, \text{ where } G \text{ is } r \times n \text{ matrix.}$$

The row rank of PA , being the same as that A , is s . Also PA has only r non-zero rows.

$\therefore$ the row rank of PA can, at the most be r

$$\therefore s \leq r \quad \dots(2)$$

From (1) and (2)

$$r = s$$

i.e. rank of A = row rank of A .

Corellary : Prove that the column rank of a matrix is the same as its rank.

Proof : We know that columns of A are the rows of A' .

$$\therefore \text{column rank of } A = \text{row rank of } A'$$

$$= \text{rank of } A'$$

$$= \text{rank of } A$$

Hence the result

Remarks :

1. Rank of A = row rank of A = column rank of A .

2. The rank of a matrix is equal to the maximum number of its linearly independent rows and also to the maximum number of its linearly independent columns.

3. If A an n -rowed non-singular matrix, then its rows as well as columns form L.I. sets.

4. If A, B be two matrices of the same type, then $\rho(A+B) \leq \rho(A) + \rho(B)$.

5. If A, B are two n-rowed square matrices, then

$$\rho(AB) \geq \rho(A) + \rho(B) - n.$$

Problem 5 : Examine the linear independent or dependence of the rows of the

matrix $A = \begin{bmatrix} 3 & 2 & 4 \\ 1 & 0 & 2 \\ 1 & -1 & -1 \end{bmatrix}$, hence find its rank.

Solution : $A = \begin{bmatrix} 3 & 2 & 4 \\ 1 & 0 & 2 \\ 1 & -1 & -1 \end{bmatrix}$

$$\therefore |A| = \begin{vmatrix} 3 & 2 & 4 \\ 1 & 0 & 2 \\ 1 & -1 & -1 \end{vmatrix} = 3 \begin{vmatrix} 0 & 2 \\ -1 & -1 \end{vmatrix} - 2 \begin{vmatrix} 1 & 2 \\ 1 & -1 \end{vmatrix} + 4 \begin{vmatrix} 1 & 0 \\ 1 & -1 \end{vmatrix}$$

$$= 3(0 + 2) - 2(-1 - 2) + 4(-1 - 0)$$

$$= 3(2) - 2(-3) + 4(-1) = 6 + 6 - 4 = 8 \neq 0$$

$\therefore$ A is non-singular matrix.

$\therefore$ three rows of A are L.I.

$\therefore$ rows of matrix A form a L.I. set

$$\therefore \rho(A) = 3.$$

Problem 6 : Find the value of k so that the vectors

$$\begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ -2 \end{bmatrix} \text{ and } \begin{bmatrix} k \\ 0 \\ 1 \end{bmatrix} \text{ are L.D.}$$

Solution : Let a, b, c be scalars, not all zero, such that

$$a \begin{bmatrix} 1 \\ -1 \\ 3 \end{bmatrix} + b \begin{bmatrix} 1 \\ 2 \\ -2 \end{bmatrix} + c \begin{bmatrix} k \\ 0 \\ 1 \end{bmatrix} = 0$$

where O is 3×1 zero matrix

$$\Rightarrow \begin{bmatrix} a \\ a \\ 3a \end{bmatrix} + \begin{bmatrix} b \\ 2b \\ -2b \end{bmatrix} + \begin{bmatrix} ck \\ 0 \\ c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow \begin{bmatrix} a + b + ck \\ -a + 2b \\ 3a - 2b + c \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\begin{aligned} \therefore \quad a + b + ck &= 0 \\ -a + 2b &= 0 \\ 3a - 2b + c &= 0 \end{aligned}$$

From (2), we get $a = 2b$

$$\therefore \quad (3) \Rightarrow 3a - a + c = 0 \Rightarrow c = -2a$$

Put the values of c and b in (1), we have

$$a + \frac{a}{2} - 2ak = 0 \Rightarrow \frac{3a}{2} - 2ak = 0 \Rightarrow 2a \left(\frac{3}{4} - k \right) = 0$$

But $a \neq 0$

[as if $a = 0$ then $b = 0$ and $c = 0$ which implies the given vectors are L.I.]

$$\Rightarrow \frac{3}{4} - k = 0 \Rightarrow k = \frac{3}{4}.$$

IV. Self check Exercise

1. Reduce to row reduced echelon form the matrix

$$A = \begin{bmatrix} 0 & 1 & 3 & -1 & 4 \\ 2 & 0 & -4 & 1 & 2 \\ 1 & 4 & 2 & 0 & -1 \\ 3 & 4 & -2 & 1 & -1 \\ 6 & 9 & -1 & 1 & 6 \end{bmatrix} \text{ and find } \rho_R(A).$$

2. Find the row rank of the matrix $A = \begin{bmatrix} 1 & 2 & -1 & 3 \\ 4 & 1 & 2 & 1 \\ 3 & -1 & 1 & 2 \\ 1 & 2 & 0 & 1 \end{bmatrix}$.

3. Show that the vectors $V_1 = (1, 2, 3)$, $V_2 = (0, 1, 2)$ and $V_3 = (0, 0, 1)$ generate $V_3(\mathbb{R})$.

4. Show that the vectors $[1 \ 2 \ 3]$, $[3 \ -2 \ -1]$, $[1 \ -6 \ -5]$ form a L.I. system.

5. Examine for linear dependence the vectors $[1, \ 2, \ 4]$, $[2, \ -1, \ 3]$, $[0, \ 1, \ 2]$, $[-3, \ 7, \ 2]$ and find the relation if it exists.

6. Prove that the vectors $x = (1, 0, 0)$, $y = (0, 1, 0)$, $z = (0, 0, 1)$ and $w = (1, 1, 1)$ form a linearly dependent set, but any three of them are linearly independent.

7. Show that the row vectors of the matrix $\begin{bmatrix} 6 & 2 & 3 & 4 \\ 0 & 5 & -3 & 1 \\ 0 & 0 & 7 & -2 \end{bmatrix}$ are linearly

independent.

8. Determine whether the following matrices have same column space or not ?

$$A = \begin{bmatrix} 1 & 3 & 5 \\ 1 & 4 & 3 \\ 1 & 1 & 9 \end{bmatrix}, B = \begin{bmatrix} 1 & 2 & 3 \\ -2 & -3 & -4 \\ 7 & 12 & 15 \end{bmatrix}.$$

EIGEN VALUES AND EIGEN VECTORS

Objectives

- I. Introduction**
- II. Some Important Results**
- III. Characteristic Equation of a Matrix**
- IV. Diagonalizable Matrix**
- V. Cayley Hamilton Theorem**
- VI. Minimal Polynomial and Minimal Equation**
- VII. Problems**
- VIII. Self Check Exercise**

I. Introduction

An expression of the form $A_0x^m + A_1x^{m-1} + A_2x^{m-2} + \dots + A_{m-1}x + A_m$, where $A_0, A_1, A_2, \dots, A_m$ are all square matrices of the same order n and m is a positive integer, is called a n -rowed matrix polynomial of degree m .

Note : Two matrix polynomials are said to be equal iff the coefficients of the like powers of x are the same.

1. Eigen values are also known as proper values, characteristic values, latent roots or spectral values. Similarly eigen vectors are also called proper vectors, characteristic vectors, latent vectors or spectral vectors.
2. The set of characteristic roots of a matrix A is called the spectrum of the matrix A .

II. Some Important Results

Result 1 : Prove that λ is an eigen value of n -rowed square matrix A over a field F if and only if $|A - \lambda I| = 0$.

Proof : (i) Assume that λ is an eigen value of A over F .

$\therefore$ there exists a non-zero column matrix X of type $n \times 1$ such that

$$AX = \lambda X$$

$$\Rightarrow AX - \lambda X = O$$

$$\Rightarrow AX - \lambda IX = O$$

$$\Rightarrow (A - \lambda I)X = O$$

$$\Rightarrow |A - \lambda I| = 0 \quad [\because X \neq O]$$

$$[\because AX = O \text{ has a non-trivial solution iff } |A - \lambda I| = 0]$$

(ii) Assume that $|A - \lambda I| = 0$

$\therefore |A - \lambda I|X = O$ has a non-trivial solution

$$\therefore AX - \lambda IX = O$$

$$\text{or } AX - \lambda X = O$$

$$\text{or } AX = \lambda X$$

Where X is a non-zero matrix

$\therefore \lambda$ is an eigen value of A over F .

Note. λ is an eigen value of A over F iff $A - \lambda I$ is a singular matrix.

Result 2 : If X is a characteristic vector of a matrix corresponding to the characteristic value λ , then kX is also a characteristic vector of A corresponding to the same characteristic value λ ($k \neq 0$).

Proof : Since X is a characteristic vector of A corresponding to the characteristic value λ .

$$\therefore X \neq O \text{ and}$$

$$AX = \lambda X$$

$$\text{Now } A(kX) = k(AX) = k(\lambda X) = \lambda(kX)$$

$$\text{Now } kX \text{ is a non-zero vector such that } A(kX) = \lambda(kX)$$

$\therefore kX$ is a characteristic vector of A corresponding to the characteristic value λ .

Note : Corresponding to a characteristic value λ , there may correspond more than one characteristic vectors.

Result 3 : If X be an eigen vector of the n -rowed square matrix A over a field F , then X cannot correspond to two distinct eigen values.

Proof : Since X is an eigen vector of A over F .

$\therefore X$ is a non-zero column matrix of order $n \times 1$.

Suppose eigen vector X corresponds to two eigen values λ_1, λ_2 of A .

$$\therefore AX = \lambda_1 X \text{ and } AX = \lambda_2 X$$

$$\Rightarrow \lambda_1 X = \lambda_2 X$$

$$\Rightarrow (\lambda_1 - \lambda_2)X = 0$$

$$\Rightarrow \lambda_1 - \lambda_2 = 0$$

Hence the result.

Result 4 : Prove that any system of eigen vectors $X_1, X_2, \dots, X_m$ corresponding respectively to a system of distinct eigen values $\lambda_1, \lambda_2, \lambda_m$ of a matrix A is linearly independent.

Proof : Try Yourself.

Result 5 : Prove that the characteristic roots of a hermitian matrix are real.

Proof : Let λ be a characteristic roots of a hermitian matrix A .

$\therefore$ there exists a non-zero $n \times 1$ column matrix X such that

$$AX = \lambda X$$

$$\Rightarrow X^0 (AX) = X^0 (\lambda X)$$

$$\Rightarrow X^0 AX = \lambda X^0 X$$

$$\Rightarrow (X^0 AX) = (\lambda X^0 X)^0$$

$$\Rightarrow X^0 A^0 (X^0)^0 = \bar{\lambda} X^0 (X^0)^0$$

$$\Rightarrow X^0 AX = \bar{\lambda} X^0 X \quad [\because A^0 = A \text{ as } A \text{ is hermitian and } (X^0)^0 = X]$$

$$\Rightarrow X^0 \lambda X = \bar{\lambda} X^0 X \quad [\because AX = \lambda X]$$

$$\Rightarrow \lambda X^0 X = \bar{\lambda} X^0 X$$

$$\Rightarrow (\lambda - \bar{\lambda}) X^0 X = 0$$

$$\Rightarrow \lambda - \bar{\lambda} = 0 \quad [\because X^0 \neq 0 \text{ as } X \neq 0]$$

$$\Rightarrow \bar{\lambda} = \lambda$$

$$\Rightarrow \lambda \text{ is real}$$

Hence the result.

Result 6 : Prove that any two characteristic vectors corresponding to two distinct characteristics roots of a hermitian matrix are orthogonal.

Proof : Let X_1, X_2 be the characteristic vectors corresponding to characteristic roots λ_1, λ_2 of the hermitian matrix A .

$$\therefore AX_1 = \lambda_1 X_1$$

$$AX_2 = \lambda_2 X_2$$

$$\text{From (1), } X_2^0 AX_1 = X_2^0 \lambda_1 X_1$$

$$\text{From (2), } X_1^0 AX_2 = X_1^0 \lambda_2 X_2$$

$$\text{Now } (X_2^0 AX_1)^0 = X_1^0 A^0 (X_2^0)^0 = X_1^0 AX_2,$$

since $A^0 = A$ as A is hermitian

$$\therefore (X_2^0 \lambda_1 X_1)^0 = X_1^0 \lambda_2 X_2$$

$$\Rightarrow \lambda_1 X_1^0 (X_2^0)^0 = \lambda_2 X_1^0 X_2 \quad [\because \lambda_1, \lambda_2 \text{ are real}]$$

$$\Rightarrow \lambda_1 X_1^0 X_2 = \lambda_2 X_1^0 X_2$$

$$\Rightarrow (\lambda_1 - \lambda_2) X_1^0 X_2 = 0$$

$$\text{But } \lambda_1 - \lambda_2 \neq 0$$

$$\therefore X_1^0 X_2 = 0$$

$$\Rightarrow X_1, X_2 \text{ are orthogonal.}$$

Result 7 : Prove that characteristic roots of a unitary matrix are of unit modulus.

Proof : Let A be given unitary matrix.

$$\therefore A^0 A = 1 \quad \dots (1)$$

Let λ be a characteristic root of A .

$\therefore$ there exists a non-zero vector X such that

$$AX = \lambda X \quad \dots (2)$$

$$\Rightarrow (AX)^\theta = (\lambda X)^\theta$$

$$\Rightarrow X^\theta A^\theta = \bar{\lambda} X^\theta \quad \dots (3)$$

From (2) and (3), we get,

$$(X^\theta A^\theta)(AX) = (\bar{\lambda} X^\theta)(\lambda X)$$

$$\Rightarrow X^\theta (A^\theta A) X = \lambda \bar{\lambda} X^\theta X$$

$$\Rightarrow X^\theta IX = \lambda \bar{\lambda} X^\theta X \quad [\because \text{of (1)}]$$

$$\Rightarrow X^\theta X = \lambda \bar{\lambda} X^\theta X$$

$$\Rightarrow (\lambda \bar{\lambda} - 1) X^\theta X = 0$$

$$\Rightarrow \lambda \bar{\lambda} - 1 = 0 \quad [\because X^\theta X \neq 0 \text{ as } X \neq 0]$$

$$\Rightarrow |\lambda|^2 - 1 = 0$$

$$\Rightarrow |\lambda|^2 = 1$$

$$\Rightarrow |\lambda| = 1$$

Hence the result.

Result 8 : Prove that any two characteristic vectors corresponding to two distinct characteristic roots of a unitary matrix are orthogonal.

Proof : Try Yourself.

III. Characteristic Equation of a Matrix

If A be any n-rowed square matrix over a field F and λ an indeterminate, then the matrix $A - \lambda I$ is called the characteristic matrix of A.

The determinant $|A - \lambda I|$, an algebraic polynomial in λ of degree n, is called the characteristic polynomial of A.

The equation $|A - \lambda I| = 0$ is called characteristic equation of A.

Remark : An eigen value λ of matrix A is always a root of its characteristic equation and every root of the characteristic equation of A is an eigen value of A.

$\therefore$ in order to find eigen values of A, we should find roots of the characteristic equation of A.

IV. Diagonalizable Matrix

An $n \times n$ matrix A is called diagonalizable if there exists an invertible $n \times n$ matrix P such that $P^{-1}AP$ is a diagonal matrix.

Method to find Diagonal Matrix for a Diagonalizable matrix.

Step I : Find eigen values $\lambda_1, \lambda_2, \dots, \lambda_n$ of A .

Step II : Find corresponding eigen vectors $X_1, X_2, \dots, X_n$. If number of eigen vectors $< n$, A is not diagonalizable.

Step III : Find $P = \{X_1 X_2 X_3 \dots X_n\}$ and P^{-1} .

Step IV : $P^{-1}AP = \text{Diag.}(\lambda_1, \lambda_2, \dots, \lambda_n)$

is required diagonal matrix.

Note : A is diagonalizable if and only if A has n L.I. eigen vectors.

V. Cayley Havilton Theorem

Statement : Every square matrix satisfies its characteristic equation.

Proof : Let A be any square matrix of order n , and its characteristic equation be

$$p_0 + p_1\lambda + p_2\lambda^2 + \dots + p_n\lambda^n = 0$$

We have to prove that A satisfies this equation

$$\text{i.e., } p_1I + p_1A + p_2\lambda^2 + \dots + p_nA^n = 0$$

For proving this, we proceed as follows :

$$\text{We know that } (A - \lambda I) \text{adj.}(A - \lambda I) = |A - \lambda I| \quad \left[\because A \text{ adj. } A = |A| I \right]$$

$$\text{Let adj. } (A - \lambda I) = B_0 + B_1\lambda + B_2\lambda^2 + \dots + B_{n-1}\lambda^{n-1}$$

$$\begin{aligned} \therefore \quad \text{we have, } (A - \lambda I)(B_0 + B_1\lambda + B_2\lambda^2 + \dots + B_{n-1}\lambda^{n-1}) \\ = (p_0 + p_1\lambda + p_2\lambda^2 + \dots + p_n\lambda^n)I \end{aligned}$$

Equating the coefficients of like powers of λ , we get,

$$AB_0 = p_0 I$$

$$AB_1 - B_0 = p_1 I$$

$$AB_2 - B_1 = p_2 I$$

$$\dots \dots \dots$$

$$AB_{n-1} - B_{n-2} = p_{n-1}I$$

$$-B_{n-1} = p_n I$$

Pre-multiplying above equations by I, A, A², ..., Aⁿ respectively and adding, we get,

$$O = p_0 I + p_1 A + p_2 A^2 + \dots + p_n A^n, \text{ which is same as (1).}$$

Hence the theorem.

VI. Minimal Polynomial and Minimal Equation

If $m(x)$ be a scalar polynomial of the lowest degree with leading coefficient unity, such that $m(x) = 0$ is satisfied by A i.e. $m(A) = O$, then the polynomial $m(x)$ is called the minimal polynomial of A and $m(x) = 0$ is called the minimum equation of A.

Note. The degree of the minimal equation of an n-rowed matrix is less than or equal to that of its characteristic equation which is n.

Derogatory and Non-derogatory Matrices

An n-rowed matrix is said to be derogatory or non-derogatory, according as the degree of its minimal equation is less than or equal to n.

VII. Problems

Problem 1 : Prove that a square matrix A and its transpose A have the same set of eigen values.

Sol. Characteristic polynomial of A^t

$$= |A^t - \lambda I| = |A - \lambda I^t| = |(A - \lambda I)^t|$$

$$= |A - \lambda I| \quad \left[\because |A^t| = |A| \right]$$

$$= \text{characteristic polynomial of A}$$

$\therefore$ A and A^t have same characteristic polynomial and hence the same set of eigen values.

Problem 2 : If α is an eigen value of a non-singular matrix A, then prove that $\frac{|A|}{\alpha}$

is an eigen value of adj. A.

Sol. Since α is an eigen of a non-singular matrix A

$\therefore \alpha \neq 0$ and there exists a non-zero column vector X such that

$$AX = \alpha X$$

$$\Rightarrow (\text{adj. A})(AX) = (\text{adj. A})(\alpha X)$$

$$\Rightarrow [(\text{adj. A})(A)]X = \alpha [(\text{adj. A})X]$$

$$\Rightarrow (|A| I) X = (\text{adj. } A) X$$

$$\Rightarrow (\text{adj. } A) X = \frac{|A|}{\alpha} X$$

$$\Rightarrow \frac{|A|}{\alpha} \text{ is an eigen value of adj. } A.$$

Problem 3 : The characteristic roots of $\begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & k \end{bmatrix}$ are 0, 3 and 15. Find the

value of k.

Sol. Let $A = \begin{bmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & k \end{bmatrix}$

$$\lambda I = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$\therefore A - \lambda I = \begin{bmatrix} 8 - \lambda & -6 & 2 \\ -6 & 7 - \lambda & -4 \\ 2 & -4 & k - \lambda \end{bmatrix}$$

$\therefore$ characteristics equation of matrix A is $|A - \lambda I| = 0$

$$\text{or } \begin{vmatrix} 8 - \lambda & -6 & 2 \\ -6 & 7 - \lambda & -4 \\ 2 & -4 & k - \lambda \end{vmatrix} = 0$$

Since $\lambda = 0$ is a root of it

$$\therefore \begin{vmatrix} 8 & -6 & 2 \\ -6 & 7 & -4 \\ 2 & -4 & k \end{vmatrix} = 0$$

$$\therefore 8 \begin{vmatrix} 7 & -4 \\ -4 & k \end{vmatrix} - (-6) \begin{vmatrix} -6 & -4 \\ 2 & k \end{vmatrix} + 2 \begin{vmatrix} -6 & 7 \\ 2 & -4 \end{vmatrix} = 0$$

$$\text{or } 8(7k - 16) + 6(-6k + 8) + 2(24 - 14) = 0$$

$$\text{or } 56k - 128 - 36k + 48 + 20 = 0$$

$$\therefore 20k = 60 \Rightarrow k = 3.$$

Problem 4 : Define similar matrices and prove that similar matrices have same characteristic polynomial and hence same eigen values.

Sol. Let A and B be square matrices of order n over a field F. Then A is said to be similar to B over F if and only if there exists an n-rowed invertible matrix P over F such that

$$AP = PB \text{ i.e. } B = P^{-1}AP \text{ or } A = PB P^{-1}$$

Let A and B be two similar matrices

$$\therefore B = P^{-1}AP$$

$$\therefore B - \lambda I = P^{-1}AP - \lambda I = P^{-1}AP - \lambda P^{-1}P \quad [\because P^{-1}P = I]$$

$$= P^{-1}AP - P^{-1}(\lambda I)P = P^{-1}(A - \lambda I)P$$

$$\therefore |B - \lambda I| = |P^{-1}(A - \lambda I)P|$$

$$= |P^{-1}| |A - \lambda I| |P|$$

$$= |A - \lambda I| |P^{-1}| |P| = |A - \lambda I| |P^{-1}P|$$

$$= |A - \lambda I| |I|$$

$$\therefore |B - \lambda I| = |A - \lambda I| \quad [\because |I| = 1]$$

$\therefore$ matrices A and B = $P^{-1}AP$ have the same characteristic polynomial and hence the same set of eigen values.

Problem 5 : Determine the eigen values and eigen vectors of the matrix

$$A = \begin{bmatrix} 3 & 1 & 1 \\ 2 & 4 & 2 \\ 1 & 1 & 3 \end{bmatrix}$$

Is it diagonalisable ? Justify.

Sol. $A = \begin{bmatrix} 3 & 1 & 1 \\ 2 & 4 & 2 \\ 1 & 1 & 3 \end{bmatrix}$

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow \lambda I = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$A - \lambda I = \begin{bmatrix} 3-\lambda & 1 & 1 \\ 2 & 4-\lambda & 2 \\ 1 & 1 & 3-\lambda \end{bmatrix}$$

$\therefore$ the characteristics equation of A is $|A - \lambda I| = 0$

or $\begin{vmatrix} 3-\lambda & 1 & 1 \\ 2 & 4-\lambda & 2 \\ 1 & 1 & 3-\lambda \end{vmatrix} = 0$

or $\begin{vmatrix} 6-\lambda & 6-\lambda & 6-\lambda \\ 2 & 4-\lambda & 2 \\ 1 & 1 & 3-\lambda \end{vmatrix} = 0$, by $R_1 \rightarrow R_1 + R_2 + R_3$

or $(6-\lambda) \begin{vmatrix} 1 & 1 & 1 \\ 2 & 4-\lambda & 2 \\ 1 & 1 & 3-\lambda \end{vmatrix} = 0$ or $(6-\lambda) \begin{vmatrix} 1 & 0 & 0 \\ 2 & 2-\lambda & 0 \\ 1 & 0 & 2-\lambda \end{vmatrix} = 0$

or $(6-\lambda)[(1)(2-\lambda)(2-\lambda)] = 0$

or $(6-\lambda)(2-\lambda)^2 = 0$

$\therefore \lambda = 2, 2, 6$

which are the eigen values of A.

The eigen vector $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \neq O$ corresponding to the eigen value $\lambda = 6$ is given by

$$AX = \lambda X \text{ or } (A - 6I)X = O$$

$$\text{or } \begin{bmatrix} -3 & 1 & 1 \\ 2 & -2 & 2 \\ 1 & 1 & -3 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \text{ or } \begin{bmatrix} 1 & 1 & -3 \\ 2 & -2 & 2 \\ -3 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 1 & 1 & -3 \\ 0 & -4 & 8 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 + R_2$$

Now the coefficient matrix of these equations is of rank 2. Therefore these equations have only $3 - 2 = 1$ L.I. solution. Thus there is only one L.I. eigen vector corresponding to the value 6. These equations can be written as

$$x + y - 3z = 0$$

$$-4y + 8z = 0 \quad \Rightarrow y = 2z$$

$$\therefore x + 2z - 3z = 0 \quad \Rightarrow x = z$$

$$\text{Take } z = 1, \quad \therefore x = 1, y = 2$$

$$X = \begin{bmatrix} 1 \\ 2 \\ 1 \end{bmatrix} \text{ is an eigen vector of A.}$$

The eigen vector $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix} \neq O$ corresponding to the eigen value $\lambda = 2$ is given by

$$AX = 2X \text{ or } (A - 2I)X = O$$

$$\text{or } \begin{bmatrix} 1 & 1 & 1 \\ 2 & 2 & 2 \\ 1 & 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{or } \begin{bmatrix} 1 & 1 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$$

The coefficient matrix of these equations is of rank 1. Therefore these equations have $3 - 1 = 2$ L.I. solutions. These equations can be written as

$$\begin{array}{ll} \mathbf{x} + \mathbf{y} + \mathbf{z} = 0 & \text{or} \quad \mathbf{x} = -\mathbf{y} - \mathbf{z} \\ \text{Take } \mathbf{y} = 1, \mathbf{z} = 0 & ; \quad \mathbf{y} = 0, \mathbf{z} = 1 \end{array}$$

Therefore we find two L.I. eigen vectors of A as $\begin{bmatrix} -1 \\ 1 \\ 0 \end{bmatrix}$ and $\begin{bmatrix} -1 \\ 0 \\ 1 \end{bmatrix}$.

$$\therefore \mathbf{P} = \begin{bmatrix} 1 & -1 & -1 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$\begin{aligned} |\mathbf{P}| &= \begin{vmatrix} 1 & -1 & -1 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{vmatrix} = 1 \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix} - (-1) \begin{vmatrix} 2 & 0 \\ 1 & 1 \end{vmatrix} + (-1) \begin{vmatrix} 2 & 1 \\ 1 & 0 \end{vmatrix} \\ &= 1(1-0) + 1(2-0) - 1(0-1) \\ &= 1(1) + 1(2) - 1(-1) = 1 + 2 + 1 \\ &= 4 \end{aligned}$$

Co-factors of the elements of first row of $|\mathbf{P}|$ are

$$= \begin{vmatrix} 1 & 0 \\ 0 & 1 \end{vmatrix}, -\begin{vmatrix} 2 & 0 \\ 1 & 1 \end{vmatrix}, \begin{vmatrix} 2 & 1 \\ 1 & 0 \end{vmatrix} \text{ i.e. } 1, 2, 1 \text{ respectively}$$

Co-factors of the elements of second row of $|\mathbf{P}|$ are

$$= \begin{vmatrix} -1 & -1 \\ 0 & 1 \end{vmatrix}, \begin{vmatrix} 1 & -1 \\ 1 & 1 \end{vmatrix}, -\begin{vmatrix} 1 & -1 \\ 1 & 0 \end{vmatrix} \text{ i.e. } 1, 2, -1 \text{ respectively}$$

Co-factors of the elements of third row of $|\mathbf{P}|$ are

$$\begin{vmatrix} -1 & -1 \\ 1 & 0 \end{vmatrix}, -\begin{vmatrix} 1 & -1 \\ 2 & 0 \end{vmatrix}, \begin{vmatrix} 1 & -1 \\ 2 & 1 \end{vmatrix} \text{ i.e. } 1, -2, 3 \text{ respectively}$$

$$\therefore \text{adj} \mathbf{P} = \begin{bmatrix} 1 & -2 & -1 \\ 1 & 2 & -1 \\ 1 & -2 & 3 \end{bmatrix}' = \begin{bmatrix} 1 & 1 & 1 \\ -2 & 2 & -2 \\ -1 & -1 & 3 \end{bmatrix}$$

$$\therefore \quad p^{-1} = \frac{\text{adj}P}{|P|} = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 \\ -2 & 2 & -2 \\ -1 & -1 & 3 \end{bmatrix}$$

$$p^{-1}AP = \frac{1}{4} \begin{bmatrix} 1 & 1 & 1 \\ -2 & 2 & -2 \\ -1 & -1 & 3 \end{bmatrix} \begin{bmatrix} 3 & 1 & 1 \\ 2 & 4 & 2 \\ 1 & 1 & 3 \end{bmatrix} \begin{bmatrix} 1 & -1 & -1 \\ 2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix}$$

$$= \frac{1}{4} \begin{bmatrix} 24 & 0 & 0 \\ 0 & 8 & 0 \\ 0 & 0 & 8 \end{bmatrix} = \begin{bmatrix} 6 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{bmatrix}$$

which is a diagonal matrix.

Problem 6 : Verify Cayley Hamilton Theorem for the matrix $A = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix}$.

Hence find A^{-1} .

Solution : $A = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix}$

$$I = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow \lambda I = \begin{bmatrix} \lambda & 0 & 0 \\ 0 & \lambda & 0 \\ 0 & 0 & \lambda \end{bmatrix}$$

$$\therefore \quad A - \lambda I = \begin{bmatrix} -\lambda & 0 & 1 \\ 1 & 2-\lambda & 0 \\ 2 & -1 & -\lambda \end{bmatrix} = -\lambda \begin{vmatrix} 2-\lambda & 0 \\ -1 & -\lambda \end{vmatrix} + 1 \begin{vmatrix} 1 & 2-\lambda \\ 2 & -1 \end{vmatrix}$$

$$= -\lambda(-2\lambda + \lambda^2) + 1(-1 - 4 + 2\lambda) = -\lambda^3 + 2\lambda^2 - 5 + 2\lambda$$

$$\therefore \quad |A - \lambda I| = -\lambda^3 + 2\lambda^2 + 2\lambda - 5$$

The characteristic equation of A is $|A - \lambda I| = 0$

$$\text{or } -\lambda^3 + 2\lambda^2 + 2\lambda - 5 = 0 \quad \text{or} \quad \lambda^3 - 2\lambda^2 - 2\lambda + 5 = 0$$

We are to prove that A satisfies this equation i.e. $A^3 - 2A^2 - 2A + 5I = O$

$$\text{Now } A^2 = \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix} = \begin{bmatrix} 2 & -1 & 0 \\ 2 & 4 & 1 \\ -1 & -2 & 2 \end{bmatrix}$$

$$A^3 = \begin{bmatrix} 2 & -1 & 0 \\ 2 & 4 & 1 \\ -1 & -2 & 2 \end{bmatrix} \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix} = \begin{bmatrix} -1 & -2 & 0 \\ 6 & 7 & 2 \\ 2 & -6 & -1 \end{bmatrix}$$

Consider $A^3 - 2A^2 - 2A + 5I$

$$= \begin{bmatrix} -1 & -2 & 2 \\ 6 & 7 & 2 \\ 2 & -6 & -1 \end{bmatrix} - 2 \begin{bmatrix} 2 & -1 & 0 \\ 2 & 4 & 1 \\ -1 & -2 & 2 \end{bmatrix} - 2 \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix} + 5 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\therefore A^3 - 2A^2 - 2A + 5I = O$$

Re-multiplying both sides by A^{-1} , we get,

$$A^2 - 2A - 2I + 5I^{-1} = 2I$$

$$= - \begin{bmatrix} 2 & -1 & 0 \\ 2 & 4 & 1 \\ -1 & -2 & 2 \end{bmatrix} + \begin{bmatrix} 0 & 0 & 1 \\ 1 & 2 & 0 \\ 2 & -1 & 0 \end{bmatrix} + 2 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$5A^{-1} = \begin{bmatrix} 0 & 1 & 2 \\ 0 & 2 & -1 \\ 5 & 0 & 0 \end{bmatrix}$$

$$\therefore A^{-1} = \frac{1}{5} \begin{bmatrix} 0 & 1 & 2 \\ 0 & 2 & -1 \\ 5 & 0 & 0 \end{bmatrix}$$

VIII. Self Check Exercise

1. If λ is an eigen value of a square matrix A, then prove that $\bar{\lambda}$ is an eigen value of A^{θ} and conversely.

2. Show that the necessary and sufficient condition for a 2×2 matrix $\begin{bmatrix} a & b \\ c & d \end{bmatrix}$ to have zero as an eigen value is that $a d - b c = 0$.

3. Determine eigen values of the matrix $\begin{bmatrix} 3 & 1 & 1 \\ 2 & 4 & 2 \\ 1 & 1 & 3 \end{bmatrix}$.

4. Find the characteristic roots and the spectrum of the matrix $\begin{bmatrix} 1 & 0 & 2 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{bmatrix}$.

5. Diagonalize, if possible, the matrix $\begin{bmatrix} 1 & 0 & 0 \\ 0 & 2 & 1 \\ 1 & -1 & 4 \end{bmatrix}$.

6. Verify Cayley-Hamilton Theorem for the matrix $A = \begin{bmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{bmatrix}$. Hence

find A^{-1} .

7. Using Cayley Hamilton theorem, find A^8 , if $A = \begin{bmatrix} 1 & 2 \\ 2 & -1 \end{bmatrix}$.

The matrix A is called the coefficient matrix.

Remarks : Any set of values $x_1, x_2, \dots, x_n$ which satisfy simultaneously the m equations in (1), is called a solution of the system.

A system of equations, which has a solution, is called consistent or compatible. If the system does not has any solution, it is called inconsistent.

II. Linearly Independent Solutions of $AX = O$

Conditions under which a set of homogeneous equations possess a (i) trivial solution of (ii) non-trivial solution.

Let there be m equations in n unknowns. So the coefficient matrix A is of type $m \times n$. Let r be rank of A.

Now either $r < n$ or $r = n$

(i) If $r = n$, then the equation $AX = O$ has $n - n = 0$

i.e. no linearly independent solution. Therefore, the equation $AX = O$ has trivial solution.

(ii) If $r < n$, then the equation $AX = O$ has $n - r$ linearly independent solutions. Any linear combination of these $n - r$ solutions will also be a solution of $AX = O$. So, there are infinite number of non-trivial solutions.

Article 1 : Let A be an $m \times n$ matrix of rank r. Then the equation $AX = O$ has $(n-r)$ linearly independent solutions.

Proof : The given equation is $AX = O \quad \dots (1)$

We have to prove two results :

(i) $AX = O$ has $(n - r)$ solutions

(ii) $(n - r)$ solutions form a linearly independent set.

For proving first part, we proceed as following :

$\therefore$ rank of A = r

$\therefore$ column rank of A = r

$\therefore$ A has r linearly independent columns. We assume that first r columns are linearly independent and last $(n-r)$ columns are linearly dependent.

Let $A = [C_1 \ C_2 \dots C_r \ C_{r+1} \dots C_n]$

$\therefore$ equation (1) can be written as

$$[C_1 \ C_2 \dots C_r \ C_{r+1} \dots C_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = O$$

$$\text{i.e., } C_1x_1 + C_2x_2 + \dots + C_rx_r + C_{r+1}x_{r+1} + \dots + C_nx_n = 0$$

$$\therefore C_{r+1}, C_{r+2}, \dots, C_n \text{ are linearly dependent columns.}$$

$$\therefore \text{each is a linear combination of } C_1, C_2, \dots, C_r$$

$$\text{Let } C_{r+1} = p_{11} C_1 + p_{12} C_2 + \dots + p_{1r} C_r$$

$$C_{r+2} = p_{21} C_1 + p_{22} C_2 + \dots + p_{2r} C_r$$

$$\dots\dots\dots$$

$$C_n = p_{t1} C_1 + p_{t2} C_2 + \dots + p_{tr} C_r \text{ where } t = n - r \quad [\because n = r + (n - r)]$$

The above equation can be written as

$$\left[\begin{array}{l} p_{11}C_1 + p_{12}C_2 + \dots + p_{1r}C_r + (-1)C_{r+1} + 0.C_{r+2} + \dots + 0.C_n = 0 \\ p_{21}C_1 + p_{22}C_2 + \dots + p_{2r}C_r + 0.C_{r+1} + (-1)C_{r+2} + \dots + 0.C_n = 0 \\ \dots\dots\dots \\ p_{t1}C_1 + p_{t2}C_2 + \dots + p_{tr}C_r + 0.C_{r+1} + 0.C_{r+2} + \dots + (-1)C_n = 0 \end{array} \right]$$

Comparing one by one the equation in (3) with equation (2), we get,

$$X_1 = \begin{bmatrix} p_{11} \\ p_{12} \\ \vdots \\ p_{1r} \\ -1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, X_2 = \begin{bmatrix} p_{21} \\ p_{22} \\ \vdots \\ p_{2r} \\ 0 \\ -1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \dots, X_t = \begin{bmatrix} p_{t1} \\ p_{t2} \\ \vdots \\ p_{tr} \\ 0 \\ 0 \\ 0 \\ \vdots \\ -1 \end{bmatrix}$$

as $t = n - r$ solutions of the equation

Now we have to show that these $n - r$ solutions $X_1, X_2, \dots, X_t$ are linearly independent vectors.

For this we consider the relation

$$p_1X_1 + p_2X_2 + \dots + p_tX_t = 0$$

$$\text{i.e. } p_1 \begin{bmatrix} p_{11} \\ p_{12} \\ \vdots \\ p_{1r} \\ -1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{bmatrix} + p_2 \begin{bmatrix} p_{21} \\ p_{22} \\ \vdots \\ p_{2r} \\ 0 \\ -1 \\ 0 \\ \vdots \\ 0 \end{bmatrix} + \dots + p_t \begin{bmatrix} p_{t1} \\ p_{t2} \\ \vdots \\ p_{tr} \\ 0 \\ 0 \\ 0 \\ \vdots \\ -1 \end{bmatrix} = 0$$

Comparing $(r + 1)$ th, $(r + 2)$ th nth elements, we get,

$$-p_1 = 0, -p_2 = 0, \dots, -p_t = 0$$

$$\therefore p_1 = p_2 = \dots = p_t = 0$$

$$\therefore p_1 X_1 + p_2 X_2 + \dots + p_t X_t = 0$$

$$\Rightarrow p_1 = p_2 = \dots = p_t = 0$$

$$\therefore X_1, X_2, \dots, X_t \text{ are L.I. vectors}$$

$$\therefore A X = 0 \text{ has } n - r \text{ L.I. solution.}$$

Article 2 : The equation $A X = 0$ has a non-zero (i.e., non-trivial solution) iff A is singular.

Proof : Assume that $A X = 0$ has a non-zero solution.

$$\therefore n - r > 0 \text{ where } r \text{ is the rank of } n\text{-rowed matrix } A \text{ implies } n > r.$$

$$\text{i.e., rank of } A \text{ is less than the order of the matrix.}$$

$$\therefore A \text{ is a singular matrix.}$$

Again, assume that A is a singular matrix

$$\therefore |A| = 0$$

$$\Rightarrow \text{rank of } A < \text{order of } A$$

$$\Rightarrow r < n$$

$$\Rightarrow n - r > 0$$

$$\Rightarrow \text{equation } A X = 0 \text{ has a non-zero solution.}$$

III. Consistency of $AX = B$

Conditions under which a system of non-homogeneous equations will have :

- (i) no solution (ii) a unique solution (iii) infinity of solutions.

Let $AX = B$ be a system of non-homogeneous equations.

- (i) The equation $AX = B$ has no solution if A and $[A \ B]$ do not have the same rank.
(ii) The equation $AX = B$, has a solution if the rank of A is the same as that of $[A \ B]$. If in addition, A is non-singular, then equation has a unique solution.
(iii) The equation $AX = B$ will have infinite solutions if A and $[A \ B]$ have the same rank and A is singular.

Article 2 : The necessary and sufficient condition that the system of equations $AX = B$ is consistent (i.e., has a solution), is that the matrices A and $[A \ B]$ are of the same rank.

Proof : Let $\rho(A) = r$ where A is $m \times n$ matrix.

$\therefore$ column rank of A is also r .

$\therefore$ r columns of A are linearly independent and the remaining $(n - r)$ are linearly dependent.

Let $C_1, C_2, \dots, C_r$ be linearly independent and $C_{r+1}, \dots, C_n$ be linearly dependent where $A = [C_1 \ C_2 \ \dots \ C_n]$.

The given equation is $Ax = B$ where $X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$

$$\text{i.e., } [C_1 \ C_2 \ \dots \ C_r \ C_{r+1} \ \dots \ C_n] \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_r \\ x_{r+1} \\ \vdots \\ x_n \end{bmatrix} = B$$

$$\text{i.e., } x_1C_1 + x_2C_2 + \dots + x_rC_r + x_{r+1}C_{r+1} + \dots + x_nC_n = B$$

Condition is necessary.

Asume that the equation $AX = B$ has a solution $X = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix}$

$$\therefore \text{ we have } \begin{bmatrix} C_1 & C_2 \dots C_r & C_{r+1} \dots C_n \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_r \\ x_{r+1} \\ \vdots \\ x_n \end{bmatrix} = B$$

$$\therefore x_1C_1 + x_2C_2 + \dots + x_rC_r + x_{r+1}C_{r+1} + \dots + x_nC_n = B \quad \dots (2)$$

$$\therefore C_{r+1}, C_{r+2}, \dots, C_n \text{ are linearly dependent and}$$

$$C_1, C_2, \dots, C_r \text{ are linearly independent.}$$

$$\therefore C_{r+1}, \dots, C_n \text{ are linear combination of } C_1, C_2, \dots, C_r \text{ and consequently from (2),}$$

B is also a linear combination of $C_1, C_2, \dots, C_r$.

$$\therefore \text{ number of linearly independent columns of } [A \quad B] \text{ is also } r.$$

$$\therefore \text{ if the equation } AX = B \text{ has a solution, then rank of } A \text{ is the same as that of } [A \quad B].$$

Condition is sufficient.

Assume rank of A as well as of $[A \quad B]$ is r .

$$\therefore \text{ rank of } [A \quad B] \text{ is } r.$$

$$\therefore \text{ number of independent columns of } [A \quad B]$$

$$\text{i.e., } [C_1 \ C_2 \dots C_r \ C_{r+1} \dots C_n \ B] \text{ is } r.$$

But $C_1, C_2, \dots, C_r$ are already linearly independent.

$$\therefore B \text{ is linearly dependent column}$$

$\therefore$ B is a linear combination of $C_1, C_2, \dots, C_r$
 $\therefore$ there exists r scalars $p_1, p_2, \dots, p_r$ such that

$$B = p_1 C_1 + p_2 C_2 + \dots + p_r C_r$$

The above equation can be written as

$$p_1 C_1 + p_2 C_2 + \dots + p_r C_r + 0 \cdot C_{r+1} + \dots + 0 \cdot C_n = B \quad \dots (3)$$

Comparing (1) and (3), we get,

$$x_1 = p_1, x_2 = p_2, \dots, x_r = p_r, x_{r+1} = \dots = x_n = 0.$$

$$\therefore X = \begin{bmatrix} p_1 \\ p_2 \\ \vdots \\ p_r \\ 0 \\ \vdots \\ 0 \end{bmatrix} \text{ is a solution of } AX = B.$$

$\therefore$ if ranks of A and $[A \ B]$ are same, the equation $AX = B$ has a solution.

Article 3 : The equation $AX = B$ has a unique solution if A is non-singular.

Proof : (i) Assume that A is non-singular i.e., A^{-1} exists.

$\therefore$ from the equation $AX = B$, we have,

$$A^{-1}(AX) = A^{-1}B \text{ i.e., } X = A^{-1}B \text{ which is a solution of } AX = B.$$

(ii) We prove that the solution is unique.

If possible, let X_1, X_2 be two different solutions of $AX = B$

$$\therefore AX_1 = B \text{ and } AX_2 = B$$

Consequently $AX_1 = AX_2$

$$\Rightarrow A^{-1}(AX_1) = A^{-1}(AX_2)$$

$$\Rightarrow X_1 = X_2$$

which is not possible as X_1, X_2 are distinct.

$\therefore$ our supposition is wrong.

$\therefore Ax = B$ has a unique solution.

IV. Problem

Problem 1 : Find the value of k so that the equation

$x - 2y + z = 0$, $3x - ty + 2z = 0$, $y + kx = 0$ have

(i) unique solution

(ii) infinitely many solution. Also find solutions for these values of k .

Solution : The given equations are

$$x - 2y + z = 0$$

$$3x - y + 2z = 0$$

$$0x + y + kz = 0$$

which can be written as

$$\begin{bmatrix} 1 & -2 & 1 \\ 3 & -1 & 2 \\ 0 & 1 & k \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore AX = O$$

$$\text{Where } A = \begin{bmatrix} 1 & -2 & 1 \\ 3 & -1 & 2 \\ 0 & 1 & k \end{bmatrix}$$

$$|A| = \begin{vmatrix} 1 & -2 & 1 \\ 3 & -1 & 2 \\ 0 & 1 & k \end{vmatrix} \xrightarrow{\text{by } R_2 \rightarrow R_2 - 3R_1} \begin{vmatrix} 1 & -2 & 1 \\ 0 & 5 & -1 \\ 0 & 1 & k \end{vmatrix}$$

$$= 1 \begin{vmatrix} 5 & -1 \\ 1 & k \end{vmatrix} = 1(5k + 1) = 5k + 1$$

(i) Equations have a unique solution

$$\text{if } |A| \neq 0$$

$$\text{i.e. if } 5k + 1 \neq 0$$

$$\text{i.e. if } k \neq -\frac{1}{5}$$

(ii) System has infinitely many solutions

$$\text{if } |A| = 0$$

$$\text{i.e. if } 5k + 1 = 0$$

i.e. if $k = -\frac{1}{5}$

When $k = -\frac{1}{5}$, we have

$$\begin{bmatrix} 1 & -2 & 1 \\ 3 & -1 & 2 \\ 0 & 1 & -\frac{1}{5} \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - 3R_1$$

$$\therefore \begin{bmatrix} 1 & -2 & 1 \\ 0 & 5 & -1 \\ 0 & 5 & -1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - R_2$$

$$\therefore \begin{bmatrix} 1 & -2 & 1 \\ 0 & 5 & -1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - R_2$$

$$\therefore x - 2y + z = 0$$

$$5y - z = 0 \Rightarrow 5y = z \Rightarrow y = \frac{1}{5}z$$

$$\therefore x - \frac{2}{5}z + z = 0 \Rightarrow x + \frac{3}{5}z = 0 \Rightarrow x = -\frac{3}{5}z$$

Put $z = k$

$$\therefore \text{ solutions are } x = -\frac{3}{5}k, y = \frac{1}{5}k, z = k, \text{ where } k \text{ is a parameter.}$$

Problem 2 : Find non-trivial solution of the system of equations

$$x - 2y - 3z = 0$$

$$-2x + 3y + 5z = 0$$

$$3x + y - 2z = 0, \text{ if possible.}$$

Sol. The given equations are

$$x - 2y - 3z = 0$$

$$-2x + 3y + 5z = 0$$

$$3x + y - 2z = 0$$

which can be written as

$$\begin{bmatrix} 1 & -2 & -3 \\ -2 & 3 & 5 \\ 3 & 1 & -2 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\therefore \begin{bmatrix} 1 & -2 & -3 \\ 0 & -1 & -1 \\ 0 & 7 & 7 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 + 2R_1, R_3 \rightarrow R_3 - 3R_1$$

$$\therefore \begin{bmatrix} 1 & -2 & -3 \\ 0 & -1 & -1 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 + 7R_2$$

$$\therefore x - 2y - 3z = 0$$

$$-y - z = 0 \Rightarrow y = -z$$

$$\therefore x + 2z - 3z = 0 \Rightarrow x = z$$

Put $z = k$

$$\therefore x = k, y = -k, z = k, \text{ where } k \text{ is a parameter.}$$

Problem 3 : Show that the system of equations

$$x + y + z = 4, 2x + 5y - 2z = 3, x + 7y - 7z = -6$$

is consistent and solve it.

Sol. The given equations are

$$x + y + z = 4$$

$$2x + 5y - 2z = 3$$

$$x + 7y - 7z = -6$$

which can be written as

$$\begin{bmatrix} 1 & 1 & 1 \\ 2 & 5 & -2 \\ 1 & 7 & -7 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ 3 \\ -6 \end{bmatrix}$$

$$\therefore \begin{bmatrix} 1 & 1 & 1 \\ 0 & 3 & -4 \\ 0 & 6 & -8 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ -5 \\ -10 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - 2R_1, R_3 \rightarrow R_3 - R_1$$

$$\therefore \begin{bmatrix} 1 & 1 & 1 \\ 0 & 3 & -4 \\ 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 4 \\ -5 \\ 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - 2R_2$$

Now rank of $\begin{bmatrix} 1 & 1 & 1 \\ 0 & 3 & -4 \\ 0 & 0 & 0 \end{bmatrix}$ as well as of $\begin{bmatrix} 1 & 1 & 1 & 4 \\ 0 & 3 & -4 & -5 \\ 0 & 0 & 0 & 0 \end{bmatrix}$ is 2.

$\therefore$ given equations are consistent and solutions are given by

$$x + y + z = 4.$$

$$3y - 4z = -5 \Rightarrow 3y = 4z - 5 \Rightarrow y = \frac{4}{3}z - \frac{5}{3}$$

$$\therefore x + \frac{4}{3}z - \frac{5}{3} + z = 4 \Rightarrow x + \frac{7}{3}z = \frac{17}{3} \Rightarrow x = -\frac{7}{3}z + \frac{17}{3}$$

Put $z = k$

$$\therefore \text{ solutions are } x = -\frac{7}{3}k + \frac{17}{3}, y = \frac{4}{3}k - \frac{5}{3}, z = k$$

where k is a parameter.

Problem 4 : Investigate for what values of a, b the following equations

$$x + y + 5z = 6$$

$$x + 2y + 3az = b$$

$$x + 3y + ax = 1$$

have

1. no solution
2. unique solution
3. an infinite number of solutions.

Sol. The given equations are

$$x + y + 5z = 6$$

$$x + 2y + 3az = b$$

$$x + 3y + az = 1$$

which can be written as

$$\begin{bmatrix} 1 & 1 & 5 \\ 1 & 2 & 3a \\ 1 & 3 & a \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ b \\ 1 \end{bmatrix}$$

$$\text{i.e., } AX = B \text{ where } A = \begin{bmatrix} 1 & 1 & 5 \\ 1 & 2 & 3a \\ 1 & 3 & a \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}, B = \begin{bmatrix} 6 \\ b \\ 1 \end{bmatrix}$$

The given equations will have a unique solution.

$$\text{if } \begin{vmatrix} 1 & 1 & 5 \\ 1 & 2 & 3a \\ 1 & 3 & a \end{vmatrix} \neq 0$$

$$\text{i.e., if } \begin{vmatrix} 1 & 1 & 5 \\ 0 & 1 & 3a-5 \\ 0 & 2 & a-5 \end{vmatrix} \neq 0, \text{ by } R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$$

$$\text{i.e., if } \begin{vmatrix} 1 & 1 & 5 \\ 0 & 1 & 3a-5 \\ 0 & 0 & -5a+5 \end{vmatrix} \neq 0, \text{ by } R_3 \rightarrow R_3 - 2R_2$$

$$\text{i.e., if } -5a+5 \neq 0 \text{ i.e., if } a \neq 1$$

$\therefore$ given equations will have a unique solution when $a \neq 1$ and b has any value
When $a = 1$,

$$A = \begin{bmatrix} 1 & 1 & 5 \\ 1 & 2 & 3 \\ 1 & 3 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 1 & 5 \\ 0 & 1 & -2 \\ 0 & 2 & -4 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$$

$$\sim \begin{bmatrix} 1 & 1 & 5 \\ 0 & 1 & -2 \\ 0 & 0 & 0 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - 2R_2$$

$$\therefore \rho(A) = 2$$

$$[A \ B] = \begin{bmatrix} 1 & 1 & 5 & 6 \\ 1 & 2 & 3 & b \\ 1 & 3 & 1 & 1 \end{bmatrix}$$

$$\sim \begin{bmatrix} 1 & 1 & 5 & 6 \\ 0 & 1 & -2 & b-6 \\ 0 & 2 & -4 & -5 \end{bmatrix}, \text{ by } R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - R_1$$

$$\sim \begin{bmatrix} 1 & 1 & 5 & 6 \\ 0 & 1 & -2 & b-6 \\ 0 & 0 & 0 & -2b+7 \end{bmatrix}, \text{ by } R_3 \rightarrow R_3 - 2R_2$$

Rank of $[A \ B]$ is 3 if $b \neq \frac{7}{2}$

$\therefore$ rank of A and $[A \ B]$ are not equal if $b \neq \frac{7}{2}$

$\therefore$ if $a = 1$, $b \neq \frac{7}{2}$, the given set of equations does not have any solution. If

$a = 1$, $b = \frac{7}{2}$, then the ranks of A and [A B] are equal and A is singular.

$\therefore$ the given system of equations has an infinite number of solutions.

V. Self Check Exercise

1. Determine the value of λ so that the equations

$$2x + y + 2z = 0$$

$$x + y + 3z = 0$$

$$4x + 3y + \lambda z = 0$$

have non-zero solution.

2. For what value of λ , does the system $\begin{bmatrix} -1 & 2 & 1 \\ 3 & -1 & 2 \\ 0 & 1 & \lambda \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = O$ has (i) a unique

solution (ii) more than one solution.

3. Solve the following equations :

$$x + y + z = 0$$

$$x + 2y + 3z = 0$$

$$x + 3y + 4z = 0$$

4. Show that the equations

$$x + y + z + 3 = 0$$

$$3x + y - 2z + 2 = 0$$

$$2x + 4y + 7z - 7 = 0$$

are inconsistent.

5. Solve the equations

$$x - y + z = 5$$

$$2x + y - z = -2$$

$$3x - y - z = -7$$

6. Examine the consistency of the following equations and if consistent, find the complete solution

$$4x - 2y + 6z = 8$$

$$x + y - 3z = -1$$

$$15x - 3y + 9z = 21$$

VECTOR SPACES-I

Objectives

- I. Vector Space (An Introduction)**
- II. Subspaces**
- III. Sum of Subspaces**
- IV. Linear Span**
- V. Self Check Exercise**

5.1 Vector Space (An Introduction)

Definition : Let $\langle V, + \rangle$ be an abelian group and $\langle F, +, \cdot \rangle$ be a field. Define a function (called scalar multiplication) from $F \times V \rightarrow V$, s.t., for all $\alpha \in F$, $v \in V$, $\alpha \cdot v \in V$. Then V is said to form a vector space over F if for all $x, y \in V$, $\alpha, \beta \in F$, the following hold

- (i) $(\alpha + \beta) x = \alpha x + \beta x$
- (ii) $\alpha (x + y) = \alpha x + \alpha y$
- (iii) $(\alpha\beta) x = \alpha (\beta x)$
- (iv) $1 \cdot x = x$, 1 being unity of F .

Also then, members of F are called scalars and those of V are called vectors.

Remark : We have used the same symbol $+$ for the two different binary compositions of V and F , for convenience. Similarly same symbol, is used for scalar multiplication and product of the field F .

Since $\langle V, + \rangle$ is group, its identity element is denoted by 0. Similarly the field F would also have zero element which will also be represented by 0.

Theorem 1 : In any vector space $V(F)$ the following results hold

- (i) $0.x = 0$
- (ii) $\alpha.0 = 0$
- (iii) $(-\alpha) x = -(\alpha x) = \alpha(-x)$
- (iv) $(\alpha - \beta)x = \alpha x - \beta x$, $\alpha, \beta \in F$, $x \in V$

Proof : (i) $0.x = (0 + 0) \cdot x = 0.x + 0.x$

$$\Rightarrow 0 + 0.x = 0.x + 0.x$$

- $\Rightarrow 0 = 0.x$ (cancellation in V)
 (ii) $\alpha.0 = \alpha.(0 + 0) = \alpha.0 + \alpha.0 \Rightarrow \alpha.0 = 0$
 (iii) $(-\alpha)x + \alpha x = [(-\alpha) + \alpha] x = 0.x = 0$
 $\Rightarrow (-\alpha x) = -\alpha x$
 (iv) follows from above.

Example 1 : Let $\langle F, +, . \rangle$ be a field

Let $V = \{(\alpha_1, \alpha_2) \mid \alpha_1, \alpha_2 \in F\}$

Define $+$ and $.$ (scalar multiplication) by

$$(\alpha_1, \alpha_2) + (\beta_1, \beta_2) = (\alpha_1 + \beta_1, \alpha_2 + \beta_2)$$

$$\alpha (\alpha_1, \alpha_2) = (\alpha\alpha_1, \alpha\alpha_2)$$

One can check that all conditions in the definition are satisfied. Here $V = F \times F = F^2$. One can extend this to F^3 and so on. In general we can take n -tuple $(\alpha_1, \alpha_2, \dots, \alpha_n)$; $\alpha_i \in F$ and define F^n or $F^{(n)} = \{(\alpha_1, \alpha_2, \dots, \alpha_n) \mid \alpha_i \in F\}$ as a Vector space over F .

Example 2 : Let $V =$ set of all real valued continuous functions defined on $[0, 1]$. Then V forms a vector space over the field R of reals under addition and scalar multiplication defined by

$$(f + g)(x) = f(x) + g(x) \quad f, g \in V$$

$$(\alpha f)(x) = \alpha f(x) \quad \alpha \in R$$

$$\text{for all } x \in [0, 1]$$

It may be recalled here that sum of two continuous functions is continuous and scalar multiple of a continuous function is continuous.

Example 3 : The set $F[x]$ of all polynomials over a field F in an indeterminate x forms a vector space over F w.r.t, the usual addition of polynomials and the scalar multiplication defined by:

$$\text{For } f(x) = a_0 + a_1x + \dots + a_nx^n \in F[x], \alpha \in F$$

$$\alpha \cdot (f(x)) = \alpha a_0 + \alpha a_1x + \dots + \alpha a_nx^n.$$

Example 4 : $M_{m \times n}(F)$, the set of all $m \times n$ matrices with entries from a field F forms a vector space under addition and scalar multiplication of matrices.

We use the notation $M_n(F)$ for $M_{n \times n}(F)$.

Example 5 : Let F be a field and X a non empty set.

Let $F^X = \{f \mid f: X \rightarrow F\}$, the set of all mappings from X to F . Then F^X forms a vector space over F under addition and scalar multiplication defined as follows:

$$\text{For } f, g \in F^X; \alpha \in F$$

Define $f + g : X \rightarrow F$, $\alpha f : X \rightarrow F$ such that

$$(f + g)(x) = f(x) + g(x)$$

$$(\alpha f)(x) = \alpha f(x) \quad \forall x \in X$$

Example 6 : Let V be the set of all vectors in three dimensional space. Addition in V is taken as the usual addition of vectors in geometry and scalar multiplication

is defined as :

$\alpha \in \mathbb{R}, \vec{v} \in V \Rightarrow \alpha \vec{v}$ is a vector in V with magnitude $|\alpha|$ times that of V . Then V forms a vector space over $\mathbb{R}$.

II. Subspaces

Definition : A non empty subset W of a vector space $V(F)$ is said to form a subspace of V if W forms a vector space under the operations of V .

Theorem 2 : A necessary and sufficient condition for a non empty subset W of a vector space $V(F)$ to be a subspace is that W is closed under addition and scalar multiplication.

Proof : If W is a subspace, the result follows by definition.

Conversely, let W be closed under addition and scalar multiplication

Let $x, y, \in W$ since $1 \in F, -1 \in F$

$$\therefore -1, y \in W \Rightarrow -y \in W$$

$$x, -y \in W \Rightarrow x - y \in W$$

$$\Rightarrow \langle W, + \rangle \text{ forms a subgroup of } \langle V, + \rangle.$$

Rest of the conditions in the definition follow trivially.

Theorem 3 : A non empty subset W of a vector space $V(F)$ is a subspace of V if $\alpha x + \beta y \in W$ for $\alpha, \beta \in F, x, y \in W$.

Proof : If W is a subspace, result follows by definition.

Conversely, let given condition hold in W .

Let $x, y \in W$ be any elements. Since $1 \in F$

$$1 \cdot x + 1 \cdot y = x + y \in W$$

$\Rightarrow W$ is closed under addition.

Again, $x \in W, \alpha \in F$ then

$$\alpha x = \alpha x + 0 \cdot y \text{ for any } y \in W, 0 \in F$$

which is in W . (Note here 0 may not be in W)

Hence W is closed under scalar multiplication.

The result thus follows by previous theorem.

Remark : V and $\{0\}$ will be trivial subspaces of any vector space $V(F)$.

Example 1 : Consider the vector space $\mathbb{R}^2(\mathbb{R})$

$$\text{then } W_1 = \{(a, 0) \mid a \in \mathbb{R}\}$$

$$W_2 = \{(0, b) \mid b \in \mathbb{R}\}$$

are subspaces of $\mathbb{R}^2$

As for any $\alpha, \beta \in \mathbb{R}, (a_1, 0), (a_2, 0) \in W_1$, we find

$$\begin{aligned} \alpha (a_1, 0) + \beta (a_2, 0) &= (\alpha a_1, 0) + (\beta a_2, 0) \\ &= (\alpha a_1 + \beta a_2, 0) \in W_1. \end{aligned}$$

Hence W_1 is a subspace. Similarly we can show W_2 is a subspace of $\mathbb{R}^2$.

Problem 1 : Show that union of two subspaces may not be a subspace.

Solution : Consider the previous example.

$W_1 \cup W_2$ will be the set containing all pairs of the type $(a, 0), (0, b)$

In particular $(1, 0), (0, 1) \in W_1 \cup W_2$

But $(1, 0) + (0, 1) = (1, 1) \notin W_1 \cup W_2$.

Hence $W_1 \cup W_2$ is not a subspace.

Reader is referred to exercises for more results pertaining to intersection and union of subspaces.

We take up few more examples of subspaces.

Problem 2 : Let $S = \{(1, 4), (0, 3)\}$ be a subset of $\mathbb{R}^2(\mathbb{R})$. Show that $(2, 3)$ belongs to $L(S)$.

Solution: $(2, 3) \in L(S)$ if it can be put as a linear combination of $(1, 4)$ and $(0, 3)$.

$$\begin{aligned}\text{Now} \quad (2, 3) &= \alpha(1, 4) + \beta(0, 3) \\ \Rightarrow (2, 3) &= (\alpha + 0, 4\alpha + 3\beta) \\ \Rightarrow 2 &= \alpha, 4\alpha + 3\beta = 3 \\ \Rightarrow \alpha &= 2, \beta = -\frac{5}{3}\end{aligned}$$

$$\text{Hence} \quad (2, 3) = 2(1, 4) - \frac{5}{3}(0, 3)$$

Showing that $(2, 3) \in L(S)$.

Problem 3 : Let $V = \mathbb{R}^4(\mathbb{R})$ and let $S = \{(2, 0, 0, 1), (-1, 0, 1, 0)\}$. Find $L(S)$.

Solution : Any element $(\alpha_1, \alpha_2, \alpha_3, \alpha_4) \in L(S)$ is a linear combination of members of S .

$$\text{Let} \quad (\alpha_1, \alpha_2, \alpha_3, \alpha_4) = \alpha(2, 0, 0, 1) + \beta(-1, 0, 1, 0), \alpha, \beta \in \mathbb{R}$$

$$\text{then} \quad (\alpha_1, \alpha_2, \alpha_3, \alpha_4) = (2\alpha - \beta, 0, \beta, \alpha)$$

$$\text{i.e.,} \quad L(S) = \{(2\alpha - \beta, 0, \beta, \alpha) \mid \alpha, \beta \in \mathbb{R}\}$$

Problem 4 : Show that the vector $F[x]$ is not finite dimensional.

Solution : Let $V = F[x]$ and suppose it is finite dimensional.

Then $\exists S \subseteq V$, s.t., $V = L(S)$ and S is finite.

Suppose $S = \{p_1, p_2, \dots, p_k\}$. We can assume $p_1 \neq 0 \quad \forall i$

Let $\deg p_i = r_i$ and let $t = \max \{r_1, r_2, \dots, r_k\}$

Now $x^{r+1} \in V$ and since $V = L(S)$,

$$x^{r+1} = \alpha_1 p_1 + \alpha_2 p_2 + \dots + \alpha_k p_k, \alpha_i \in F$$

$$\text{So} \quad 0 = (-1) x^{t+1} + \alpha_1 p_1 + \dots + \alpha_k p_k$$

Since x^{r+1} does not appear in $p_1, p_2, \dots, p_k$

we get $-1 = 0$, a contradiction. Hence V is not FDVS over F .

Note if $S = \{1, x, \dots, x^n, \dots\}$ then $V = L(S)$.

Example 2 : Let $V = R[x]$ and suppose $W = \{f(x) \in V \mid f(x) = f(1-x)\}$

Then W is a subspace of V as

$W \neq \phi$ since $0 \in W$ as $f(x) = 0 = f(1-x)$

Again, if $f(x), g(x) \in W$, then $f(x) = f(1-x), g(x) = g(1-x)$

Let $f(x) + g(x) = h(x)$

Then $h(1-x) = f(1-x) + g(1-x)$

$$= f(x) + g(x) = h(x)$$

$\Rightarrow h(x) \in W$ or that $f(x) + g(x) \in W$

Again, for $\alpha \in R$, let $\alpha f(x) = r(x)$

Then $r(1-x) = \alpha f(1-x) = \alpha f(x) = r(x)$

$\Rightarrow r(x) \in W \Rightarrow \alpha f(x) \in W$

Hence W is a subspace.

Example 3 : Let $V = F^X$ (see example 7) and suppose $Y \subseteq X$

Then $W = \{f \in V \mid f(y) = 0 \quad \forall y \in Y\}$ is a subspace of V

Clearly $0 \in W$ and for $f, g \in W$, $f(y) = 0 = g(y) \quad \forall y \in Y$

So $(f+g)(y) = f(y) + g(y) = 0 \quad \forall y \in Y$

$$\Rightarrow f+g \in W$$

Again, if $\alpha \in F$, then $(\alpha f)(y) = \alpha(f(y)) = 0 \quad \forall y \in Y$

$$\Rightarrow \alpha f \in W.$$

Example 4 : If $V = R^n$, then

$W = \{x_1, x_2, \dots, x_n \mid x_1 + x_2 + \dots + x_n = 1\}$ will not be subspace of V .

Notice, $(1, 0, 0, \dots, 0) + (0, 1, 0, \dots, 0) = (1, 1, 0, \dots, 0) \notin W$.

Example 5 : Let $V = M_{2 \times 1}(F)$. Let A be a 2×2 matrix over F .

Then $W = \left\{ \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in V \mid A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0 \right\}$ forms a subspace of V

$$W \neq \phi \text{ as } \begin{bmatrix} 0 \\ 0 \end{bmatrix} \in W$$

For $\begin{bmatrix} x_1 \\ x_2 \end{bmatrix}, \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$ in W , we have

$$A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0 = A \begin{bmatrix} y_1 \\ y_2 \end{bmatrix}$$

$$\Rightarrow A \left(\begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \right) = 0$$

$$\Rightarrow \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \end{bmatrix} \in W$$

Also $A \left(\alpha \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \right) = \alpha A \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = 0 \Rightarrow \alpha \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} \in W$

Hence W is a subspace of V .

III. Sum of Subspaces

If W_1 and W_2 be two subspaces of a vector space $V(F)$ then we define

$$W_1 + W_2 = \{w_1 + w_2 \mid w_1 \in W_1, w_2 \in W_2\}$$

$$W_1 + W_2 \neq \phi \text{ as } 0 = 0 + 0 \in W_1 + W_2$$

Again, $x, y \in W_1 + W_2$, $\alpha, \beta \in F$ implies

$$x = w_1 + w_2$$

$$y = w'_1 + w'_2 \quad w_1, w'_1 \in W_1, w_2, w'_2 \in W_2$$

$$\alpha x + \beta y = \alpha(w_1 + w_2) + \beta(w'_1 + w'_2)$$

$$= (\alpha w_1 + \beta w'_1) + (\alpha w_2 + \beta w'_2) \in W_1 + W_2$$

Showing thereby that sum of two subspaces is a subspace.

One can extend the definition, similarly, to the sum of n subspaces $W_1, W_2, \dots, W_n$,

which would also be a subspace and we write $W_1 + W_2 + \dots + W_n = \sum_{i=1}^n W_i$.

Definition : Let $W_1, W_2, \dots, W_n$ be subspaces of V then $W_1 + W_2 + \dots + W_n$ is called the direct sum if each $x \in W_1 + W_2 + \dots + W_n$ can be expressed uniquely as $x = w_1 + w_2 + \dots + w_n, w_i \in W_i$ and in that case we write

$$W_1 + W_2 + \dots + W_n = W_1 \oplus W_2 \oplus \dots \oplus W_n$$

We say, a vector space V is the direct sum of its subspaces $W_1, W_2, \dots, W_n$ if $V = W_1 \oplus W_2 \oplus \dots \oplus W_n$, i.e., if

$$V = W_1 + W_2 + \dots + W_n$$

and each $v \in V$ can be expressed uniquely as $v = w_1 + w_2 + \dots + w_n, w_i \in W_i$.

Theorem 4 : $V = W_1 \oplus W_2 \Leftrightarrow V = W_1 + W_2, W_1 \cap W_2 = \{0\}$.

Proof : Let $V = W_1 \oplus W_2$

We need to prove $W_1 \cap W_2 = \{0\}$

Let $x \in W_1 \cap W_2$, then $x \in W_1$ and $x \in W_2$

$$\Rightarrow x = 0 + x \in W_1 + W_2 = V$$

$$\Rightarrow x = x + 0 \in W_1 + W_2 = V$$

Since x has been expressed as $x = x + 0$ and $0 + x$ and the representation has to be unique, we get $x = 0$

$$\Rightarrow W_1 \cap W_2 = \{0\}.$$

Conversely, let $v \in V$ be any element and suppose

$$v = w_1 + w_2$$

$$v = w'_1 + w'_2$$

are two representation of v

$$\text{then } w_1 + w_2 = w'_1 + w'_2 (= v)$$

$$\Rightarrow w_1 - w'_1 = w'_2 - w_2$$

Now L.H.S. is in W_1 and R.H.S. belongs to W_2

i.e., each belongs to $W_1 \cap W_2 = \{0\}$

$$\Rightarrow w_1 = w'_1 = w'_2 - w_2 = 0$$

$$\Rightarrow w_1 = w'_1, w_2 = w'_2.$$

Hence the result.

Remark: The above theorem can also be stated as

$$W_1 + W_2 = W_1 \oplus W_2 \Leftrightarrow W_1 \cap W_2 = \{0\}.$$

If W be a subspace of a vector space $V(F)$ then since $\langle W, + \rangle$ forms an abelian group

of $\langle V, +, \rangle$, we can talk of cosets of W in V . Let $\frac{V}{W}$ be the set of all cosets $W + v$, $v \in V$,

then we show that $\frac{V}{W}$ also forms a vector space over F , under the operations defined by

$$\begin{aligned}(W + x) + (W + y) &= W + (x + y) & x, y \in V \\ \alpha(W + x) &= W + \alpha x & \alpha \in F\end{aligned}$$

Addition is well defined, since,

$$\begin{aligned}W + x &= W + x' \\ W + y &= W + y' \\ \Rightarrow x - x' &\in W, y - y' \in W \\ \Rightarrow (x - x') + (y - y') &\in W \\ \Rightarrow (x + y) - (x' + y') &\in W \\ \Rightarrow W + (x + y) &= W + (x' + y')\end{aligned}$$

Again,

$$\begin{aligned}W + x &= W + x' \\ \Rightarrow x - x' &\in W \\ \Rightarrow \alpha(x - x') &\in W & \alpha \in F \\ \Rightarrow \alpha x - \alpha x' &\in W \\ \Rightarrow W + \alpha x &= W + \alpha x' \\ \Rightarrow \alpha(W + x) &= \alpha(W + x')\end{aligned}$$

Thus, scalar multiplication is also well defined. It should now be a routine exercise to check that all conditions in the definition of a vector space are satisfied.

$W + 0$ will be zero of $\frac{V}{W}$

$W - x$ will be inverse of $W + x$

IV. Linear Span

Definition : Let $V(F)$ be a vector space, $v_i \in V, \alpha_i \in F$ be elements of V and F

respectively. Then elements of the type $\sum_{i=1}^n \alpha_i v_i$ are called linear combinations of

$v_1, v_2, \dots, v_n$ over F .

Let S be a non empty subset of V , then the set

$$L(S) = \left\{ \sum_{i=1}^n \alpha_i v_i \mid \alpha_i \in F, v_i \in S, n \text{ finite} \right\}$$

i.e., the set of all linear combinations of finite sets of elements of S is called linear span of S . It is also denoted by $\langle S \rangle$. If $S = \emptyset$, define $L(S) = \{0\}$.

Theorem 5 : $L(S)$ is the smallest subspace of V , containing S .

Proof : $L(S) \neq \emptyset$ as $v \in S \Rightarrow v = 1 \cdot v, 1 \in F$

$$\Rightarrow v \in L(S)$$

thus, in fact, $S \subseteq L(S)$.

Let $x, y \in L(S), \alpha, \beta \in F$ be any elements

then $x = \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n$

$$y = \beta_1 v'_1 + \beta_2 v'_2 + \dots + \beta_m v'_m \quad v_i, v'_j \in S, \alpha_i, \beta_j \in F$$

$$\text{Thus } \alpha x + \beta y = \alpha \alpha_1 v_1 + \alpha \alpha_2 v_2 + \dots + \alpha \alpha_n v_n + \beta \beta_1 v'_1 + \dots + \beta \beta_m v'_m.$$

R.H.S. being a linear combination belongs to $L(S)$

Hence $L(S)$ is a subspace of V , containing S .

Let now W be any subspace of V , containing S

We show $L(S) \subseteq W$

$$x \in L(S) \Rightarrow x = \sum \alpha_i v_i \quad v_i \in S, \alpha_i \in F$$

$v_i \in S \subseteq W$ for all i and W is a subspace

$$\Rightarrow \sum \alpha_i v_i \in W \Rightarrow x \in W$$

$$\Rightarrow L(S) \subseteq W$$

Hence the result follows.

Theorem 6 : If W is a subspace of V , then $L(W) = W$ and conversely.

Proof : $W \subseteq L(W)$ by definition and since $L(W)$ is the smallest subspace of V containing W and W is itself a subspace

$$L(W) \subseteq W$$

$$\text{Hence } L(W) = W.$$

Conversely, let $L(W) = W$

$$\text{Let } x, y \in W, \alpha, \beta \in F$$

$$\text{Then } x, y \in L(W)$$

- $\Rightarrow$ x, y are linear combinations of members of W .
- $\Rightarrow$ $\alpha x + \beta y$ is a linear combination of members of W
- $\Rightarrow$ $\alpha x + \beta y \in L(W)$
- $\Rightarrow$ $\alpha x + \beta y \in W$
- $\Rightarrow$ W is a subspace.

Definition : If $V = L(S)$, we say S spans (or generates) V . The vector space V is said to be finite-dimensional (over F) if there exists a finite subset S of V such that $V = L(S)$. We use notation F.D.V.S. for a finite dimensional vector space.

It now follows from the results we've proved that

If S_1 and S_2 are two subspaces of V , then $S_1 + S_2$ is the subspace spanned by $S_1 \cup S_2$

Indeed, $L(S_1 \cup S_2) = L(S_1) + L(S_2) = S_1 + S_2$.

V. Self Check Exercise :

1. Let $\alpha_1 = (1, 1, -2, 1)$, $\alpha_2 = (3, 0, 4, -1)$, $\alpha_3 = (-1, 2, 5, 2)$. Show that the vector $(4, -5, 9, -7)$ is spanned by $\alpha_1, \alpha_2, \alpha_3$.
2. If S spans V then show that every super set of S spans V .

VECTOR SPACES-II

I. Linear Dependence and Independence

II. Basis and Dimensions

III. Self Check Exercise

I. Linear Dependence and Independence

Let $V(F)$ be a vector space. Elements $v_1, v_2, \dots, v_n$ in V are said to be linearly dependent (over F) if $\exists$ scalars $\alpha_1, \alpha_2, \dots, \alpha_n \in F$, (not all zero) such that

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0$$

($v_1, v_2, \dots, v_n$ are finite in number, not essentially distinct).

Thus for linear dependence $\sum \alpha_i v_i = 0$ and at least one $\alpha_i \neq 0$.

If $v_1, v_2, \dots, v_n$ are not linearly dependent (L.D.) these are called linearly independent (L.I.).

In other words, $v_1, v_2, \dots, v_n$ are L.I. if

$$\sum \alpha_i v_i = 0 \Rightarrow \alpha_i = 0 \text{ for all } i$$

A finite set $X = \{x_1, x_2, \dots, x_n\}$ is said to be L.D. or L.I. according as its n members are L.D. or L.I.

In general any subset Y of $V(F)$ is called L.I. if every finite non empty subset of Y is L.I. otherwise it is called L.D.

So, if some subsets are L.I. and some are L.D. then Y is called L.D.

Observations : (i) A non zero vector is always L.I. as $v \neq 0, \alpha v = 0$ would mean $\alpha = 0$.

(ii) Zero vector is always L.D.

$$1. 0 = 0 \quad 1 \neq 0, 1 \in F$$

Thus any collection of vectors to which zero belongs is always L.D.

In other words, if $v_1, v_2, \dots, v_n$ are L.I. then none of these can be zero.

(iii) v is L.I. iff $v \neq 0$.

(iv) Any subset of a L.I. set is L.I.

- (v) Any super set of a L.D. set is L.D.
 (vi) Empty set ϕ is L.I. since it has no non empty finite subset and consequently it satisfies the condition for linear independence.
 (vii) A set of vector is L.I. if and only if every finite subset of it is L.I.

Example 1 : Consider $R^2(R)$, $R = \text{reals}$.

$$v_1 = (1, 0), v_2 = (0, 1) \in R^2 \text{ are L.I.}$$

as
$$\alpha_1 v_1 + \alpha_2 v_2 = 0 \text{ for } \alpha_1, \alpha_2 \in R$$

$$\Rightarrow \alpha_1 (1, 0) + \alpha_2 (0, 1) = (0, 0)$$

$$\Rightarrow (\alpha_1, \alpha_2) = (0, 0) \Rightarrow \alpha_1 = \alpha_2 = 0.$$

Example 2 : Consider the subset

$$S = \{(1, 0, 0), (0, 1, 0), (0, 0, 1), (2, 3, 4)\}$$

in the vector space $R^3(R)$

Since $2(1, 0, 0) + 3(0, 1, 0) + 4(0, 0, 1) - 1(2, 3, 4) = (0, 0, 0)$
 we find S is L.D.

Example 3 : In the vector space $F[x]$ of polynomials the vectors $f(x) = 1 - x$, $g(x) = x - x^2$, $h(x) = 1 - x^2$ are L.D. since $f(x) + g(x) - h(x) = 0$.

Problem 1 : Show that the vectors $v_1 = (0, 1, -2)$, $v_2 = (1, -1, 1)$, $v_3 = (1, 2, 1)$ are L.I. in $R^3(R)$.

Solution : Let $\sum \alpha_i v_i = 0$ for $\alpha_i \in R$

Then
$$\alpha_1 (0, 1, -2) + \alpha_2 (1, -1, 1) + \alpha_3 (1, 2, 1) = (0, 0, 0)$$

$$\Rightarrow (0, \alpha_1, -2\alpha_1) + (\alpha_2, -\alpha_2, \alpha_2) + (\alpha_3, 2\alpha_3, \alpha_3) = (0, 0, 0)$$

$$\Rightarrow 0 + \alpha_2 + \alpha_3 = 0$$

$$\alpha_1 - \alpha_2 + 2\alpha_3 = 0$$

$$-2\alpha_1 + \alpha_2 + \alpha_3 = 0$$

Since the coefficient determinant $\begin{vmatrix} 0 & 1 & 1 \\ 1 & -1 & 2 \\ -2 & 1 & 1 \end{vmatrix}$ is $-6 \neq 0$ the above equations have

only the zero common solution

$$\Rightarrow \alpha_1 = \alpha_2 = \alpha_3 = 0 \Rightarrow v_1, v_2, v_3 \text{ are L.I.}$$

Problem 2 : Show that $\{f(x), g(x), h(x)\}$ is L.I. in $F[x]$, whenever. $\deg f(x)$, $\deg g(x)$, $\deg h(x)$ are distinct.

Solution : Let $f(x) = a_0 + a_1x + \dots + a_mx^m, a_m \neq 0$

$$g(x) = b_0 + b_1x + \dots + b_nx^n, b_n \neq 0$$

$$h(x) = c_0 + c_1x + \dots + c_tx^t, c_t \neq 0$$

$$\text{Let } \alpha f(x) + \beta g(x) + \gamma h(x) = 0, \alpha, \beta, \gamma \in F$$

Let $m < n < t$ (without any loss of generality)

$$\text{then } \gamma C_t = 0 \Rightarrow \gamma = 0 \text{ as } c_t \neq 0$$

$$\therefore \alpha f(x) + \beta g(x) = 0$$

$$\text{and so } \beta b_n = 0 \Rightarrow \beta = 0 \text{ as } b_n \neq 0$$

$$\Rightarrow \alpha f(x) = 0 \Rightarrow \alpha a_m = 0 \Rightarrow \alpha = 0 \text{ as } a_m \neq 0$$

Hence $\{f(x), g(x), h(x)\}$ is L.I. in $F[x]$ over F .

Problem 3 : Show that the vectors

$v_1 = (1, 1, 2, 4)$, $v_2 = (2, -1, -5, 2)$, $v_3 = (1, -1, -4, 0)$ and $v_4 = (2, 1, 1, 6)$ are L.D. in $R^4(R)$.

Solution : Suppose $av_1 + bv_2 + cv_3 + dv_4 = 0, a, b, c, d \in R$

$$\text{then } a(1, 1, 2, 4) + b(2, -1, -5, 2)$$

$$+ c(1, -1, -4, 0) + d(2, 1, 1, 6) = (0, 0, 0, 0)$$

$$\text{or } (a, a, 2a, 4a) + (2b, -b, -5b, 2b) + (c, -c, -4c, 0) + (2d, d, d, 6d) = (0, 0, 0, 0)$$

$$\Rightarrow a + 2b + c + 2d = 0$$

$$a - b - c + d = 0$$

$$2a - 5b - 4c + d = 0$$

$$4a + 2b + 0c + 6d = 0$$

$$\Rightarrow \begin{bmatrix} 1 & 2 & 1 & 2 \\ 1 & -1 & -1 & 1 \\ 2 & -5 & -4 & 1 \\ 4 & 2 & 0 & 6 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$R_2 \rightarrow R_2 - R_1, R_3 \rightarrow R_3 - 2R_1, R_4 \rightarrow R_4 - 4R_1$$

$$\begin{bmatrix} 1 & 2 & 1 & 2 \\ 0 & -3 & -2 & -1 \\ 0 & -3 & -2 & -1 \\ 0 & -3 & -2 & -1 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$R_4 \rightarrow \frac{1}{2}R_4, R_3 \rightarrow \frac{1}{3}R_3$$

$$\begin{bmatrix} 1 & 2 & 1 & 2 \\ 0 & -3 & -2 & -1 \\ 0 & -1 & -2/3 & -1/3 \\ 0 & -3/4 & -1 & -1/2 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$R_4 \rightarrow R_4 - R_2, R_3 \rightarrow R_3 - R_2$$

$$\begin{bmatrix} 1 & 2 & 1 & 2 \\ 0 & -3 & -2 & -1 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} a \\ b \\ c \\ d \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\Rightarrow a + 2b + c + 2d = 0$$

$$-3b - 2c + d = 0$$

$$3b + 2c + d = 0$$

$a = -1, b = -1, c = 1, d = 1$ satisfy the equations.

Since coefficients are non zero, the given vectors are L.D.

Problem 4 : Show that

(i) $\{1, \sqrt{2}\}$ is L.I in R over Q .

(ii) $\{1, \sqrt{2}, \sqrt{3}\}$ is L.I in R over Q .

(iii) $\{1, \sqrt{2}, \sqrt{3}, \sqrt{6}\}$ is L.I in R over Q .

Solution : (i) Suppose $a + b\sqrt{2} = 0, a, b \in Q$

Suppose $b \neq 0$, then $\sqrt{2} = -\frac{a}{b} \in Q$, a contradiction

Hence $b = 0$ and so $a = 0$. Thus $\{1, \sqrt{2}\}$ is L.I. in R over Q .

(ii) Let $a + b\sqrt{2} + c\sqrt{3} = 0$, $a, b, c \in Q$

Let $c \neq 0$, then

$$\sqrt{3} = -\frac{a}{c} - \frac{b}{c}\sqrt{2} = \alpha + \beta\sqrt{2}, \alpha, \beta \in Q$$

$$\Rightarrow 3 = \alpha^2 + \alpha\beta^2 + 2\alpha\beta\sqrt{2}$$

$$\Rightarrow \alpha\beta\sqrt{2} \in Q \Rightarrow \alpha\beta = 0$$

Let $\alpha = 0$ then $\beta = \sqrt{\frac{3}{2}}$, a contradiction

So, $c = 0$ giving $a + b\sqrt{3} = 0 \Rightarrow a = b = 0$ by (i)

Hence the result follows.

(iii) Let $a + b\sqrt{2} + c\sqrt{3} + d\sqrt{6} = 0$, $a, b, c, d \in Q$

$$\text{Then } (a + b\sqrt{2}) + \sqrt{3}(c + d\sqrt{2}) = 0$$

Let $c + d\sqrt{2} \neq 0$

$$\text{Then } \sqrt{3} = \frac{-(a + b\sqrt{2})}{(c + d\sqrt{2})} = \frac{-(a + b\sqrt{2})(c - d\sqrt{2})}{c^2 - 2d^2}$$

$$= \alpha + \beta\sqrt{2}, \quad \alpha, \beta \in Q$$

$$\Rightarrow \alpha.1 + \beta\sqrt{2} + (-1)\sqrt{3} = 0$$

$$\Rightarrow -1 = 0 \text{ by (ii), a contradiction}$$

$$\therefore c + d\sqrt{2} = 0 \Rightarrow c = d = 0 \Rightarrow a + b\sqrt{2} = 0$$

$$\Rightarrow a = b = 0$$

Hence the result follows.

Problem 5 : If two vectors are L.D. then one of them is scalar multiple of the other.

Solution : Suppose v_1, v_2 are L.D. then $\exists \alpha_1 \in F$, s.t.,

$$\alpha_1 v_1 + \alpha_2 v_2 = 0 \text{ for some } \alpha_1 \neq 0$$

without loss of generality we can take $\alpha_1 \neq 0$, then α_1^{-1} exists and $\alpha_1 v_1 = (-\alpha_2 v_2)$

$$\Rightarrow v_1 = (-\alpha_1^{-1} \alpha_2) v_2 = \beta v_2$$

which proves the result.

Problem 6 : If x, y, z are L.I. over the field C of complex nos. then so are $x + y + z$ and $z + x$ over C .

Solution : Suppose $\alpha_1(x + y) + \alpha_2(y + z) + \alpha_3(z + x) = 0, \alpha_i \in C$

$$\text{Then } (\alpha_1 + \alpha_3)x + (\alpha_1 + \alpha_2)y + (\alpha_2 + \alpha_3)z = 0$$

$$\alpha_1 + \alpha_3 = \alpha_1 + \alpha_2 = \alpha_2 + \alpha_3 = 0 \text{ as } x, y, z \text{ are L.I.}$$

Solving we find

$$\alpha_1 = \alpha_2 = \alpha_3 = 0$$

Hence the result.

II. Basis and Dimensions

Note : We have already showed that $(1, 0)$ and $(0, 1)$ are L.I. in $R^2(R)$. If

$v = (a, b) \in R^2$ be any element then since $(a, b) = a(1, 0) + b(0, 1)$, $a, b \in R$

We find any element of R^2 can be written as a linear combination of $\{(1, 0), (0, 1)\} = S$

$$\text{i.e., } v \in R^2 \Rightarrow v \in L(S)$$

$$\Rightarrow R^2 \subseteq L(S)$$

$$\text{But } L(S) \subseteq R^2$$

$$\text{i.e., } R^2 \subseteq L(S)$$

or the S spans R^2 .

Definition : Let $V(F)$ be a vector space. A subset S of V is called a basis of V if S consists of L.I. elements (i.e., any finite number of elements in S are L.I.) and $V = L(S)$, i.e., S spans V .

Therefore, $S = \{(1, 0), (0, 1)\}$ is a basis of $R^2(R)$. It is rather easy to see then that $\{(1, 0), (0, 1), (0, 0, 1)\}$ will form basis of $R^3(R)$, and one can trivially extend this to $R^{(n)}$.

Again $\{(1, 1, 0), (1, 0, 0), (0, 1, 1)\}$ also forms a basis of $\mathbb{R}^3(\mathbb{R})$. (Show !) Thus a vector space may have more than one basis.

If the elements in a basis are written in a certain specific order, we call it ordered basis. Also $\{(1, 0), (0, 1)\}$, $\{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$ etc. are called standard basis of $\mathbb{R}^2$, $\mathbb{R}^3$ etc. Also ϕ is a basis for $V = \{0\}$.

Problem 7 : Show that the set $S = \{(1, 2, 1), (2, 1, 0), (1, -1, 2)\}$ form a basis of $\mathbb{R}^3(\mathbb{R})$.

Solution : Let

$$\alpha_1(1, 2, 1) + \alpha_2(2, 1, 0) + \alpha_3(1, -1, 2) = (0, 0, 0), \alpha_i \in \mathbb{R}$$

$$\Rightarrow (\alpha_1 + 2\alpha_2 + \alpha_3, 2\alpha_1 + \alpha_2 - \alpha_3, \alpha_1 + 0 + 2\alpha_3) = (0, 0, 0)$$

$$\Rightarrow \alpha_1 + 2\alpha_2 + \alpha_3 = 0$$

$$2\alpha_1 + \alpha_2 - \alpha_3 = 0$$

$$\alpha_1 + 0 + 2\alpha_3 = 0$$

In matrix form, we get
$$\begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -1 \\ 1 & 0 & 2 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

$$\text{i.e. } AX = 0$$

$$\text{where } |A| = -9 \neq 0$$

So A is a non singular matrix and thus $AX = 0$ has the unique zero solution

$$\alpha_1 = \alpha_2 = \alpha_3 = 0.$$

Hence S is L.I. set

Again, to show that $L(S) = \mathbb{R}^3$, let $(a, b, c) \in \mathbb{R}^3$ be any element. We want that $(a, b, c) = \beta_1(1, 2, 1) + \beta_2(2, 1, 0) + \beta_3(1, -1, 2)$ for same $\beta_1, \beta_2, \beta_3 \in \mathbb{R}$

i.e., we want some $\beta_1 \in \mathbb{R}$ s.t., the equations

$$\beta_1 + 2\beta_2 + \beta_3 = a$$

$$2\beta_1 + \beta_2 - \beta_3 = b$$

$$\beta_1 + 0\beta_2 - 2\beta_3 = c$$

are satisfied i.e, in matrix form

$$AX = B \text{ where } A = \begin{bmatrix} 1 & 2 & 1 \\ 2 & 1 & -1 \\ 1 & 0 & 2 \end{bmatrix}, B = \begin{bmatrix} a \\ b \\ c \end{bmatrix}$$

Since $|A| = -9 \neq 0$, $AX = B$ has a unique solution i.e., $\exists$ some β_i s.t., above equation are satisfied or that it is possible to express any $(a, b, c) \in \mathbb{R}^3$ as a linear combination members of S . i.e., $L(S) = \mathbb{R}^3$

Hence S forms a basis of $\mathbb{R}^3(\mathbb{R})$.

Theorem 1 : If $S = \{v_1, v_2, \dots, v_n\}$ is basis of V , thenf every element of V can be expressed uniquely as a linear combination of $v_1, v_2, \dots, v_n$.

Proof : Since, by definition of basis, $V = L(S)$, each element $v \in V$ can be expressed as linear combination of $v_1, v_2, \dots, v_n$.

$$\text{Suppose } v = \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n, \alpha_i \in F$$

$$v = \beta_1 v_1 + \beta_2 v_2 + \dots + \beta_n v_n, \beta_i \in F$$

$$\text{then } \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = \beta_1 v_1 + \beta_2 v_2 + \dots + \beta_n v_n$$

$$\Rightarrow (\alpha_1 - \beta_1)v_1 + (\alpha_2 - \beta_2)v_2 + \dots + (\alpha_n - \beta_n)v_n = 0$$

$$\Rightarrow \alpha_i - \beta_i = 0 \text{ for all } i (v_1, v_2, \dots, v_n \text{ are L.I.})$$

$$\Rightarrow \alpha_i = \beta_i \text{ for all } i.$$

Theorme 2 : Suppose S is a finite subset of a vector space V such that $V = L(S)$ [i.e., V is F.D.V.S] then there exists a subset of S which is a basis of V .

Proof : If S consists of L.I. elements then S itself forms basis of V and we've nothing to prove.

Let now T be a subset of S , such that T spans V and T is such minimal subset of S . (Existence of T is ensured as S is finite).

$$\text{Suppose } T = \{v_1, v_2, \dots, v_n\}$$

we show T is L.I.

$$\text{Let } \sum \alpha_i v_i = 0, \alpha_i \in F$$

Suppose $\alpha_i \neq 0$ for some i . Without any loss of generality we can take $\alpha_1 \neq 0$.

Then α_1^{-1} exists.

$$\text{Now } \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0$$

$$\Rightarrow \alpha_1^{-1} (\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n) = 0$$

$$\Rightarrow v_1 = (-\alpha_1^{-1}\alpha_2)v_2 + (-\alpha_1^{-1}\alpha_3)v_3 + \dots + (-\alpha_1^{-1}\alpha_n)v_n$$

$$= \beta_2 v_2 + \beta_3 v_3 + \dots + \beta_n v_n \quad \beta_i \in F$$

If $v \in V$ be any element then

$$v = \gamma_1 v_1 + \gamma_2 v_2 + \dots + \gamma_n v_n \quad \gamma_i \in F \quad \text{as } V = L(T)$$

$$\Rightarrow v = \gamma_1 (\beta_2 v_2 + \dots + \beta_n v_n) + \gamma_2 v_2 + \dots + \gamma_n v_n$$

i.e., any element of V is a linear combination of $v_2, v_3, \dots, v_n$

$\Rightarrow \{v_2, v_3, \dots, v_n\}$ spans V , which contradicts our choice of T (as T was such minimal)

Hence $\alpha_1 = 0$

or that $\alpha_i = 0$ for all i

$$\Rightarrow v_1, v_2, \dots, v_n \text{ are L.I.}$$

and thus T is a basis V .

Cor : A F.D.V.S has a basis.

In fact, one can prove this result for any vector space (i.e. any vector space has a basis).

Theorem 3 : If V is a F.D.V.S and $\{v_1, v_2, \dots, v_r\}$ is a L.I. subset of V , then it can be extended to form a basis of V .

Proof : If $\{v_1, v_2, \dots, v_r\}$ spans V , then it self forms a basis of V and there is nothing to prove.

Let $S = \{v_1, v_2, \dots, v_r, v_{r+1}, \dots, v_n\}$ be the maximal L.I. subset of V , containing $\{v_1, v_2, \dots, v_r\}$.

We show S is a basis of V , for which it is enough to prove that S spans V .

Let $v \in V$ be any element

then $T = \{v_1, v_2, \dots, v_n, v\}$ is L.D. by choice of S

$\Rightarrow \exists \alpha_1, \alpha_2, \dots, \alpha_n, \alpha \in F$ (not all zero) such that

$$\alpha_1 v_1 + \dots + \alpha_n v_n + \alpha v = 0$$

We claim $\alpha \neq 0$. Suppose $\alpha = 0$

then $\alpha_1 v_1 + \dots + \alpha_n v_n = 0$

$\Rightarrow \alpha_i = 0$ for all i as $v_1, v_2, \dots, v_n$ are L.I.

$\therefore \alpha = \alpha_i = 0$ for all i which is not true.

Hence $\alpha \neq 0$ and so α^{-1} exists.

Since $v = (\alpha^{-1}\alpha_1)v_1 + (\alpha^{-1}\alpha_2)v_2 + \dots + (\alpha^{-1}\alpha_n)v_n$

v is a linear combination of $v_1, v_2, \dots, v_n$

which proves our assertion.

Theorem 4 : If $\dim V = n$ and $S = \{v_1, v_2, \dots, v_n\}$ spans V then S is a basis of V .

Proof : Since $\dim V = n$, any basis of V has n elements. But theorem 8, a subset of S will be a basis of V but as S contains n elements, it will itself form basis of V .

Theorem 5 : If $\dim V = n$ and $S = \{v_1, v_2, \dots, v_n\}$ is L.I. subset of V then S is a basis of V .

Proof : Since $\{v_1, v_2, \dots, v_n\} = S$ is L.I. it can be extended to form a basis of V , but $\dim V$ being n it will itself be a basis of V .

Aliter : Let $v \in V$, then

$v, v_1, v_2, \dots, v_n$ will be L.D.. Thus $\exists \alpha, \alpha_1, \alpha_2, \dots, \alpha_n \in F$ s.t.,

$$\alpha v + \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0$$

where some α_i or α is not zero.

If $\alpha = 0$, then

$$\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n = 0$$

$$\Rightarrow \alpha_i = 0 \forall i \text{ as } v_1, v_2, \dots, v_n \text{ are L.I.}$$

Thus $\alpha \neq 0$ and so

$$v = (-\alpha^{-1}\alpha_1)v_1 + \dots + (-\alpha^{-1}\alpha_n)v_n \in L(S)$$

$$\Rightarrow V \subseteq L(S)$$

$$\Rightarrow V = L(S) \text{ and as } S \text{ is L.I. } S \text{ is a basis of } V.$$

Problem 8 : If $\{v_1, v_2, \dots, v_n\}$ is a basis of F.D.V.S V of $\dim n$ and $v =$

$\sum \alpha_i v_i, \alpha_r \neq 0$ then prove that $\{v_1, v_2, \dots, v_{r-1}, v, v_{r+1}, \dots, v_n\}$ is also a basis of V .

Solution : We have

$$v = \alpha_1 v_1 + \dots + \alpha_r v_r + \dots + \alpha_n v_n \quad \alpha_r \neq 0 \therefore \alpha_r^{-1} \text{ exists}$$

$$\Rightarrow v_1 = (\alpha_r^{-1} \alpha_1) v_1 + \dots + (-\alpha_r^{-1} \alpha_{r-1}) v_{r-1} + \alpha_r^{-1} v + \dots + (-\alpha_r^{-1} \alpha_n) v_n$$

$$\Rightarrow \beta_1 v_1 + \dots + \beta_{r-1} v_{r-1} + \beta_r v + \beta_{r+1} + \dots + \beta_n v_n$$

If $x \in V$ be any element, then

$$x = \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n \quad \alpha_i \in F$$

$$\Rightarrow x = \alpha_1 v_1 + \dots + \alpha_{r-1} v_{r-1} + \alpha_r (\beta_1 v_1 + \dots + \beta_n v_n) + \dots + \alpha_n v_n$$

or that x is a linear combination of

$$v_1, \dots, v_{r-1}, v, v_{r+1}, \dots, v_n$$

and x being any element, we find V is spanned by $\{v_1, \dots, v_{r-1}, v, v_{r+1}, \dots, v_n\}$ and it forms a basis of V , using theorem done above.

Theorem 6 : Two finite dimensional vector spaces over F are isomorphic iff they have the same dimension.

Proof : Let V and W be two isomorphic vector spaces over F and let $\theta : V \rightarrow W$ be the isomorphism.

Let $\dim V = n$ and $\{v_1, v_2, \dots, v_n\}$ be a basis of V .

We claim $\{\theta(v_1), \theta(v_2), \dots, \theta(v_n)\}$ is basis of W .

$$\text{Now } \sum_{i=1}^n \alpha_i \theta(v_i) = 0 \quad \alpha_i \in F$$

$$\Rightarrow \sum \theta(\alpha_i v_i) = 0 = \theta(0)$$

$$\Rightarrow \sum \alpha_i v_i = 0 \quad (\theta \text{ is } 1-1)$$

$$\Rightarrow \alpha_i = 0 \text{ for all } i \text{ as } v_1, v_2, \dots, v_n \text{ are L.I.}$$

$$\Rightarrow \theta(v_1), \theta(v_2), \dots, \theta(v_n) \text{ are L.I.}$$

Again, we $w \in W$ is any element, then as θ is onto, $\exists$ some $v \in V$ s.t., $\theta(v) = w$

$$\text{Now } v \in V \Rightarrow v = \sum_{i=1}^n \alpha_i v_i \text{ for some } \alpha_i \in F$$

$$\Rightarrow w = \theta(v) = \theta\left(\sum \alpha_i v_i\right)$$

$$\Rightarrow 'w = \sum \theta(\alpha_i v_i) = \alpha_1 \theta(v_1) + \alpha_2 \theta(v_2) + \dots + \alpha_n \theta(v_n)$$

or that w is a linear combination of $\theta(v_1), \theta(v_2), \dots, \theta(v_n)$

Hence $\theta(v_1), \theta(v_2), \dots, \theta(v_n)$ span W and therefore, form a basis of W showing that $\dim W = n$.

Conversely, let $\dim V = \dim W = n$ and suppose, $\{v_1, v_2, \dots, v_n\}$ and $\{w_1, w_2, \dots, w_n\}$ are basis of V and W respectively.

Define a map $\theta : V \rightarrow W$ s.t.,

$$\theta(v) = \theta(\alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n)$$

$$\alpha_1 w_1 + \alpha_2 w_2 + \dots + \alpha_n w_n$$

then θ is easily seen to be well defined. (Indeed any $v \in V$ is unique linear combination of members of basis).

If $v, v' \in V$ be any elements then

$$v = \sum \alpha_i v_i, v' = \sum \beta_i v_i \quad \alpha_i, \beta_i \in F$$

$$\begin{aligned} \theta(v + v') &= \theta\left(\sum \alpha_i v_i + \sum \beta_i v_i\right) \\ &= \theta\left(\sum (\alpha_i + \beta_i) v_i\right) \\ &= \sum (\alpha_i + \beta_i) w_i \\ &= \sum \alpha_i w_i + \sum \beta_i w_i = \theta(v) + \theta(v') \end{aligned}$$

$$\text{Also } \theta(\alpha v) = \theta\left(\alpha \sum \alpha_i v_i\right) = \theta\left(\sum \alpha \alpha_i v_i\right) = \sum (\alpha \alpha_i) w_i$$

$$\alpha \sum \alpha_i w_i = \alpha \theta(v)$$

Thus θ is a homomorphism

Now if $v \in \text{Ker } \theta$

then $\theta(v) = 0$

$$\Rightarrow \theta\left(\sum \alpha_i v_i\right) = 0$$

$$\Rightarrow \sum \alpha_i w_i = 0$$

$$\Rightarrow \alpha_i = 0 \text{ for all } i \text{ } w_1, w_2, \dots, w_n \text{ being L.I.}$$

$$\Rightarrow v = 0$$

$$\Rightarrow \text{Ker } \theta = \{0\}$$

$\Rightarrow \theta$ is one-one.

That θ is onto is obvious. Hence θ is an isomorphism.

Problem 9 : $(1, 1, 1)$ is L.I. vector in $\mathbb{R}^3(\mathbb{R})$ Extend it to form a basis of $\mathbb{R}^3$.

Solution : $(1, 1, 1)$ is non zero vector and it therefore L.I. in $\mathbb{R}^3$.

Let $S = \{(1, 1, 1)\}$, then $L(S) = \{\alpha(1, 1, 1) \mid \alpha \in \mathbb{R}\}$

Now $(1, 0, 0) \in \mathbb{R}^3$, but $(1, 0, 0) \notin L(S)$

thus by above problem $S_1 = \{(1, 1, 1), (1, 0, 0)\}$ is L.I.

Now $L(S_1) = \{\alpha(1, 1, 1) + \beta(1, 0, 0) \mid \alpha, \beta \in \mathbb{R}\}$

$$= \{(\alpha + \beta, \alpha, -\alpha) \mid \alpha, \beta \in \mathbb{R}\}$$

Again $(0, 1, 0) \notin L(S_1)$ and by above problem

$S_2 = \{(1, 1, 1), (1, 0, 0), (0, 1, 0)\}$ is L.I. subset of $\mathbb{R}^3$.

Since $\dim \mathbb{R}^3 = 3$, we find S_2 will be a basis of $\mathbb{R}^3$.

Theorem 7 : Let W be a subspace of a F.D.V.S. V , then W is finite dimensional and $\dim W \leq \dim V$. In fact, $\dim V = \dim W$ iff $V = W$.

Proof : Let $\dim V = n$, then n is maximum number of L.I. elements in any subset of V . Since any subset of W will be a subset of V , n is the maximum number of L.I. elements in W .

Let $w_1, w_2, \dots, w_m$ be the maximum number of L.I. elements in W when $m \leq n$.

We show $\{w_1, w_2, \dots, w_m\}$ is basis of W . These are already L.I. If $w \in W$ be any element then the set $\{w_1, w_2, \dots, w_m, w\}$ is L.D.

$\Rightarrow \exists \alpha_1, \alpha_2, \dots, \alpha_m, \alpha$ in F (not all zero) s.t.,

$$\alpha_1 w_1 + \dots + \alpha_m w_m + \alpha w = 0$$

If $\alpha = 0$ we get $\alpha_i = 0$ for all i as $w_1, \dots, w_m$ are L.I. which is not true. Thus $\alpha \neq 0$ and so α^{-1} exists.

The above equation then gives us

$$w = (-\alpha^{-1}\alpha_1)w_1 + \dots + (-\alpha^{-1}\alpha_m)w_m$$

Showing that $\{w_1, w_2, \dots, w_m\}$ spans W (and thus W is finite dimensional)

$\Rightarrow \{w_1, w_2, \dots, w_m\}$ is a basis of W

$$\Rightarrow \dim W = m \leq n = \dim V$$

Finally, if $\dim V = \dim W = n$

and $\{w_1, w_2, \dots, w_m\}$ be a basis of W then as $\{w_1, w_2, \dots, w_n\}$ is L.I. in W it will be L.I. in V . and as $\dim V = n$, $\{w_1, w_2, \dots, w_n\}$ is a basis of V .

Now if $v \in V$ be any element then

$$\Rightarrow v = \alpha_1 w_1 + \alpha_2 w_2 + \dots + \alpha_n w_n \in W$$

$$\Rightarrow V \subseteq W \Rightarrow V = W$$

Conversely, of course, $V = W \Rightarrow \dim V = \dim W$.

Theorem 8 : Let W be a subspace of a F.D.V.S. V . Then

$$\dim \frac{V}{W} = \dim V - \dim W.$$

Proof : Let $\dim W = m$ and let $\{w_1, w_2, \dots, w_m\}$ be a basis of W .

$w_1, w_2, \dots, w_m$ being L.I. in W will be L.I. in V and thus $\{w_1, w_2, \dots, w_m\}$ can be extended to form a basis of V .

Let $\{w_1, w_2, \dots, w_m, v_1, v_2, \dots, v_n\}$ be this extended basis of V .

Then $\dim V = n + m$

Consider the set $S = \{w + v_1, w + v_2, \dots, w + v_n\}$, we show it forms a basis of $\frac{V}{W}$.

$$\text{Let } \alpha_1 (w + v_1) + \dots + \alpha_n (w + v_n) = w, \alpha_i \in F$$

$$\text{Then } w + (\alpha_1 v_1 + \dots + \alpha_n v_n) = w$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n \in W$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n \text{ is a linear combination of } w_1, \dots, w_m$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n = \beta_1 w_1 + \dots + \beta_m w_m, \beta_j \in F$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n - \beta_1 w_1 - \dots - \beta_m w_m = 0$$

$$\Rightarrow \alpha_i = \beta_j = 0 \text{ for all } i, j.$$

$$\Rightarrow \{w + v_1, w + v_2, \dots, w + v_n\} \text{ is L.I.}$$

Again, for any $w + v \in \frac{V}{W}$, $v \in V$ means v is a linear combination of $w_1, \dots, w_m, v_1, \dots, v_n$.

$$\text{i.e., } v = \alpha_1 w_1 + \dots + \alpha_m w_m + \beta_1 v_1 + \dots + \beta_n v_n, \alpha_i, \beta_j \in F$$

$$\begin{aligned}
\text{giving } W + v &= W + (\alpha_1 w_1 + \dots + \alpha_m w_m) + (\beta_1 v_1 + \dots + \beta_n v_n) \\
&= W + (\beta_1 v_1 + \dots + \beta_n v_n) \\
&= (W + \beta_1 v_1) + \dots + (W + \beta_n v_n) \\
&= \beta_1 (W + v_1) + \beta_2 (W + v_2) + \dots + \beta_n (W + v_n).
\end{aligned}$$

Hence S spans $\frac{V}{W}$ and is therefore a basis.

$$\therefore \dim \frac{V}{W} = n$$

$$\text{Thus } \dim \frac{V}{W} = \dim V - \dim W.$$

Theorem 9 : If A and B are two subspaces of a F.D.V.S. V then

$$\dim (A + B) = \dim A + \dim B - \dim (A \cap B).$$

Proof : We've already proved that

$$\frac{A+B}{A} \cong \frac{B}{A \cap B}$$

$$\therefore \dim \frac{A+B}{A} = \dim \frac{B}{A \cap B}$$

$$\Rightarrow \dim (A+B) - \dim A = \dim B - \dim (A \cap B)$$

$$\text{or that } \dim (A+B) = \dim A + \dim B - \dim (A \cap B).$$

Remark : The reader should try to give an independent proof of the above theorem as an exercise.

Cor. : If $A \cap B = (0)$ then $\dim (A + B) = \dim A + \dim B$

$$\text{i.e., } \dim (A \oplus B) = \dim A + \dim B.$$

III. Self Check Exercise

1. Show that the following vectors are L.I.
 - (i) $(1, 0, 0), (1, 1, 1), (1, 2, 3)$, in $\mathbb{R}^3(\mathbb{R})$
 - (ii) $(1, 2, -1), (2, 2, 1), (1, -2, 3)$ in $\mathbb{R}^3(\mathbb{R})$
2. Show that the following vectors are L.D.
 - (i) $(1, 1, 2), (-3, 1, 0), (1, -1, 1), (1, 2, -3)$ in $\mathbb{R}^3(\mathbb{R})$

- (ii) $(1, 1, 2), (1, 2, 5), (5, 3, 4)$ in $R^3(R)$
3. Show that $\{1, i\}$ forms a basis of $C(R)$.
 4. Extend the set $S = \{(1, 1, 0)\}$ to form two different bases of $R^3(R)$.
 5. Let S be a finite subset of a vector space V such that S is L.I. and every proper superset of S in V is L.D. Show that S is a basis of V .
 6. If W_1 and W_2 are subspaces of R^4 and $\{(1, 0, 0, 0), (1, 1, 0, 0), (1, 1, 1, 0)\}, \{(0, 0, 0, 1), (0, 0, 1, 1), (0, 1, 1, 1)\}$ are bases of W_1 and W_2 respectively, find a basis of $W_1 \cap W_2$.
 7. Let V be a vector space over F . Assume that every linearly independent set in V can be extended to a basis of V . Deduce that V has a basis.
 8. Let F be a field. Let $A = \{(x, y, 0) \mid x, y, \in F\}$, $B = \{(0, y, z) \mid y, z \in F\}$ be subspaces of $F^3(F)$. Find dimension of the subspace $A + B$.

LINEAR TRANSFORMATIONS-I

Objectives

- I. Linear Transformations (An Introduction)**
- II. Rank and Nullity of a Linear Transformation**
- III. Self Check Exercise**

I. Linear Transformations (An Introduction)

Definition : Let V and U be two vector spaces over the same field F , then a mapping $T : V \rightarrow U$ is called a homomorphism or a linear transformation if

$$T(x + y) = T(x) + T(y) \text{ for all } x, y \in V$$

$$T(\alpha x) = \alpha T(x) \quad \alpha \in F$$

One can combine the two conditions to get a single condition

$$T(\alpha x + \beta y) = \alpha T(x) + \beta T(y) \quad x, y \in V; \alpha, \beta \in F$$

It is easy to see that both are equivalent. If a homomorphism happens to be one-one onto also we call it an isomorphism, and say the two spaces are isomorphic. (Notation $V \cong U$).

Example 1 : Identity map $I : V \rightarrow V$, s.t.,

$$I(v) = v$$

and the zero map $O : V \rightarrow V$, s.t.,

$$O(v) = 0$$

are clearly linear transformations.

Example 2 : For a field F , consider the vector spaces F^2 and F^3 . Define a map $T : F^3 \rightarrow F^2$, by

$$T(\alpha, \beta, \gamma) = (\alpha, \beta)$$

then T is a linear transformation as

for any $x, y \in F^3$, if $x = (\alpha_1, \beta_1, \gamma_1)$

$$y = (\alpha_2, \beta_2, \gamma_2)$$

$$\begin{aligned} \text{then } T(\mathbf{x} + \mathbf{y}) &= T(\alpha_1 + \alpha_2, \beta_1 + \beta_2, \gamma_1 + \gamma_2) = (\alpha_1 + \alpha_2, \beta_1 + \beta_2) \\ &= (\alpha_1, \beta_1) + (\alpha_2, \beta_2) = T(\mathbf{x}) + T(\mathbf{y}) \end{aligned}$$

$$\begin{aligned} \text{and } T(\alpha \mathbf{x}) &= T(\alpha(\alpha_1, \beta_1, \gamma_1)) = T(\alpha\alpha_1, \alpha\beta_1, \alpha\gamma_1) \\ &= (\alpha\alpha_1, \alpha\beta_1) = \alpha(\alpha_1, \beta_1) = \alpha T(\mathbf{x}) \end{aligned}$$

Example 3 : Let V be the vector space of all polynomials in x over a field F . Define $T : V \rightarrow V$, s.t.,

$$T(f(x)) = \frac{d}{dx} f(x)$$

$$\text{then } T(f + g) = \frac{d}{dx}(f + g) = \frac{d}{dx}f + \frac{d}{dx}g = T(f) + T(g)$$

$$T(\alpha f) = \frac{d}{dx}(\alpha f) = \alpha \frac{d}{dx}f = \alpha T(f)$$

shows that T is a linear transformation.

In fact if $\theta : V \rightarrow V$ be defined such that

$$\theta(f) = \int_0^x f(t) dt$$

then θ will also be a linear transformation.

Example 4 : Consider the mapping

$$T : \mathbb{R}^3 \rightarrow \mathbb{R}, \text{ s.t.,}$$

$$T(x_1, x_2, x_3) = x_1^2 + x_2^2 + x_3^2$$

then T is not a linear transformation.

Consider, for instance,

$$T((1, 0, 0) + (1, 0, 0)) = T(2, 0, 0) = 4$$

$$T(1, 0, 0) + T(1, 0, 0) = 1 + 1 = 2.$$

In the following theorems, we take V and U to be vector spaces over the same field.

Theorem 1 : Under a homomorphism $T : V \rightarrow U$,

$$(i) T(0) = 0 \quad (ii) T(-x) = -T(x).$$

Proof : $T(0) = T(0 + 0) = T(0) + T(0)$

$$\Rightarrow T(0) = 0$$

$$\text{Again } T(-x) + T(x) = T(-x + x) = T(0) = 0$$

$$\Rightarrow -T(x) = T(-x).$$

Definition : Let $T : V \rightarrow U$ be a homomorphism, then kernel of T is defined by

$$\text{Ker } T = \{x \in V \mid T(x) = 0\}$$

It is also called the null space of T .

Theorem 2 : Let $T : V \rightarrow U$ be a homomorphism, then $\text{Ker } T$ is a subspace of V .

Proof : $\text{Ker } T \neq \emptyset$, as $0 \in \text{Ker } T$

Let $\alpha, \beta \in F$, $x, y \in \text{Ker } T$ be any elements

$$\begin{aligned} \text{then } T(\alpha x + \beta y) &= \alpha T(x) + \beta T(y) \\ &= \alpha \cdot 0 + \beta \cdot 0 = 0 + 0 = 0 \end{aligned}$$

$$\Rightarrow \alpha x + \beta y \in \text{Ker } T.$$

Theorem 3 : Let $T : V \rightarrow U$ be a homomorphism, then

$$\text{Ker } T = \{0\} \text{ iff } T \text{ is one-one.}$$

Proof : Let $\text{Ker } T = \{0\}$. If $T(x) = T(y)$

$$\text{then } T(x) - T(y) = 0$$

$$\Rightarrow T(x - y) = 0$$

$$\Rightarrow (x - y) \in \text{Ker } T = \{0\}$$

$$\Rightarrow x - y = 0$$

$$\Rightarrow x = y \Rightarrow T \text{ is 1-1.}$$

Conversely, let T be one-one

if $x \in \text{Ker } T$ be any element, then $T(x) = 0$

$$\Rightarrow T(x) = T(0)$$

$$\Rightarrow x = 0$$

$$\Rightarrow \text{Ker } T = \{0\}.$$

Definition : Let $T : V \rightarrow U$ be a linear transformation then range of T is defined to

be

$$\begin{aligned} T(V) &= \{T(x) \mid x \in V\} = \text{Range } T = R_T \\ &= \{u \in U \mid u = T(v), v \in V\} \end{aligned}$$

Theorem 4 : Let $T : V \rightarrow U$ be a L.T. (linear transformation) then range of T is subspace of U .

Proof : Since $T(0) = 0, 0 \in V$

$$\therefore T(0) \in \text{Range } T$$

$$\text{i.e., } \text{Range } T \neq \emptyset$$

Let $\alpha, \beta \in F, T(x), T(y) \in T(V)$ be any elements

$$\text{then } x, y \in V$$

$$\text{Now } \alpha T(x) + \beta T(y) = T(\alpha x + \beta y) \in T(V)$$

$$\text{as } \alpha x + \beta y \in V$$

Hence the result.

Note: $T(V) = U$ iff T is onto.

Theorem 5 : Let $T : V \rightarrow U$ be a L.T. then

$$\frac{V}{\text{Ker } T} \cong \text{Range } T = T(V).$$

Proof : Let $T : V \rightarrow U$ and put $\text{Ker } T = K$, then K being a subspace of V , we can talk V/K .

Define a mapping $\theta : V/K \rightarrow T(V)$, s.t.,

$$\theta(K + x) = T(x), x \in V$$

Then θ is well defined, one-one map as

$$K + x = K + y$$

$$\Leftrightarrow x - y \in K = \text{Ker } T$$

$$\Leftrightarrow T(x - y) = 0$$

$$\Leftrightarrow T(x) = T(y)$$

$$\Leftrightarrow \theta(K + x) = \theta(K + y)$$

If $T(x) \in T(V)$ be any element, then $x \in V$ and $\theta(K + x) = T(x)$, showing that θ is onto

Finally, $\theta((K + x) + (K + y)) = \theta(K + (x + y))$

$$= T(x + y)$$

$$= T(x) + T(y)$$

$$= \theta(K + x) + \theta(K + y)$$

$$\text{and } \theta(\alpha(K + x)) = \theta(K + \alpha x) = T(\alpha x) = \alpha T(x) = \alpha \theta(K + x)$$

shows θ is a L.T. and hence an isomorphism.

Note : The above is called the Fundamental Theorem of homomorphism for vector

space. If the map T is also onto, then we have proved $\frac{V}{\text{Ker } T} \cong U$.

Theorem 6 : If A and B be two subspaces of a vector space $V(F)$, then

$$\frac{A+B}{A} \cong \frac{B}{A \cap B}.$$

Proof : A being a subspace of $A + B$ and $A \cap B$ being a subspace of B , we can talk

of $\frac{A+B}{A}$ and $\frac{B}{A \cap B}$.

Define a map $\theta: B \rightarrow \frac{A+B}{A}$, s.t.,

$$\theta(b) = A + b, b \in B$$

Since $b_1 = b_2 \Rightarrow A + b_1 = A + b_2$, we find θ is well defined.

Again, as $\theta(\alpha b_1 + \beta b_2) = A + (\alpha b_1 + \beta b_2)$

$$= (A + \alpha b_1) + (A + \beta b_2)$$

$$= \alpha(A + b_1) + \beta(A + b_2)$$

$$= \alpha\theta(b_1) + \beta\theta(b_2)$$

θ is a L.T.

For any $A + x \in \frac{A+B}{A}$, we find $x \in A + B$

$$\Rightarrow x = a + b, a \in A, b \in B$$

$$\begin{aligned}
A + x &= A + (a + b) \\
&= (A + a) + (A + b) = A + (A + b) \\
&= A + b = \theta(b).
\end{aligned}$$

Showing that b is the required pre image of $A + x$ under θ and thus θ is onto. Hence by Fundamental theorem

$$\frac{A+B}{A} \cong \frac{B}{\text{Ker } \theta}.$$

We claim $\text{Ker } \theta = A \cap B$

$$\begin{aligned}
\text{Indeed } x \in \text{Ker } \theta &\Leftrightarrow \theta(x) = A \\
&\Leftrightarrow A + x = A \\
&\Leftrightarrow x \in A, \text{ also } x \in \text{Ker } \theta \subseteq B \\
&\Leftrightarrow x \in A \cap B
\end{aligned}$$

$$\text{Hence } \frac{A+B}{A} \cong \frac{B}{A \cap B}$$

Note : By interchanging A and B , we get $\frac{B+A}{B} \cong \frac{A}{B \cap A}$

$$\text{i.e., } \frac{A+B}{A} \cong \frac{B}{A \cap B}.$$

Cor.: If $A + B$ is the direct sum then as $A \cap B = \{0\}$

$$\text{we get } \frac{A}{(0)} \cong \frac{A \oplus B}{B}$$

$$\text{But } \frac{A}{(0)} \cong A \text{ gives us } A \cong \frac{A \oplus B}{B}.$$

Theorem 7 : Let W be a subspace of V , then $\exists$ an onto L.T. $\theta: V \rightarrow \frac{V}{W}$ such that $\text{Ker } \theta = W$.

Proof : The proof is left as an exercise for the reader.

Theorem 8 : A L.T. $T : V \rightarrow V$ is one-one iff T is onto.

Proof : Let $T : V \rightarrow V$ be one-one. Let $\dim V = n$.

Let $\{v_1, v_2, \dots, v_n\}$ be a basis of V , then $\{T(v_1), \dots, T(v_n)\}$ will also be a basis of V as

$$\alpha_1 T(v_1) + \alpha_2 T(v_2) + \dots + \alpha_n T(v_n) = 0$$

$$\Rightarrow T(\alpha_1 v_1 + \dots + \alpha_n v_n) = T(0) \quad (T \text{ a L.T.})$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n = 0 \quad (T \text{ is 1-1})$$

$$\Rightarrow \alpha_i = 0 \text{ for all } i$$

thus $T(v_1), \dots, T(v_n)$ are L.I. and as $\dim V = n$ the result follows (Theorem done earlier)

Let now $v \in V$ be any element

$$\text{then} \quad v = a_1 T(v_1) + a_2 T(v_2) + \dots + a_n T(v_n) \quad a_i \in F$$

$$= T(a_1 v_1 + \dots + a_n v_n)$$

$$= T(v') \text{ for some } v'$$

Hence T is onto.

Conversely, let T be onto.

Here again we show that if $\{v_1, v_2, \dots, v_n\}$ is a basis of V then so also is

$$\{T(v_1), T(v_2), \dots, T(v_n)\}$$

For any $v \in V$, since T is onto, $\exists$ some $v' \in V$ s.t.,

$$T(v') = v$$

II. Rank and Nullity of a Linear Transformation

Definition : Let $T : V \rightarrow W$ be a L.T.

then we define Rank of $T = \dim \text{Range } T = r(T)$

Nullity of $T = \dim \text{Ker } T = v(T)$.

Theorem 9 : (Sylvester's Law) : Let $T : V \rightarrow W$ be a L.T., then

$$\text{Rank } T + \text{Nullity } T = \dim V.$$

Proof : Let $\{x_1, x_2, \dots, x_m\}$ be a basis of $\text{Ker } T$ then $\{x_1, x_2, \dots, x_m\}$ being L.I. in $\text{Ker } T$ will be L.I. in V . Thus it can be extended to form a basis of V .

Let $\{x_1, x_2, \dots, x_m, v_1, v_2, \dots, v_n\}$ be the extended basis of V .

Then $\dim \text{Ker } T = \text{nullity of } T = m$
 $\dim V = m + n$

we show $\{T(v_1), T(v_2), \dots, T(v_n)\}$ is a basis of $\text{Range } T$

Now $\alpha_1 T(v_1) + \alpha_2 T(v_2) + \dots + \alpha_n T(v_n) = 0$

$$\Rightarrow T(\alpha_1 v_1 + \dots + \alpha_n v_n) = 0$$

$$\Rightarrow \alpha_1 v_1 + \alpha_2 v_2 + \dots + \alpha_n v_n \in \text{Ker } T$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n = \beta_1 x_1 + \dots + \beta_m x_m$$

or $\alpha_1 v_1 + \dots + \alpha_n v_n + (-\beta_1)x_1 + \dots + (-\beta_m)x_m = 0$

$$\Rightarrow \alpha_1 = \alpha_2 = \dots = \beta_1 = \dots = \beta_m = 0$$

i.e., $\{T(v_1), T(v_2), \dots, T(v_n)\}$ is L.I.

Now if $T(v) \in \text{Range } T$ be any element then as $v \in V$

$$v = a_1 x_1 + \dots + a_m x_m + b_1 v_1 + \dots + b_n v_n \quad a_i, b_j \in F$$

$$T(v) = a_1 T(x_1) + \dots + a_m T(x_m) + b_1 T(v_1) + \dots + b_n T(v_n)$$

$$= 0 + \dots + 0 + b_1 T(v_1) + \dots + b_n T(v_n) \quad [\text{as } x_i \in \text{Ker } T]$$

or that $T(v)$ is a linear combination of $T(v_1), \dots, T(v_n)$

which, therefore, form a basis of $\text{Range } T$.

$\therefore \dim \text{Range } T = n = \text{rank } T$

which proves the theorem.

Theorem 10 : If $T: V \rightarrow V$ be a L.T. Show that the following statements are equivalent.

- (i) $\text{Range } T \cap \text{Ker } T = \{0\}$
- (ii) If $T(T(v)) = 0$ then $T(v) = 0, v \in V$

Proof : (i) $\Rightarrow$ (ii)

$$T(T(v)) = 0 \Rightarrow T(v) \in \text{Ker } T$$

Also $T(v) \in \text{Range } T$ (by definition)

$$\Rightarrow T(v) = 0$$

(ii) $\Rightarrow$ (i)

Let $x \in \text{Range } T \cap \text{Ker } T$

$$\Rightarrow x \in \text{Range } T \text{ and } x \in \text{Ker } T$$

$$\Rightarrow x = T(v) \text{ for some } v \in V$$

$$\text{and } T(x) = 0$$

$$x = T(v) \Rightarrow T(x) = T(T(v))$$

$$\Rightarrow 0 = T(T(v))$$

$$\Rightarrow T(v) = 0 \text{ (given condition)}$$

$$\Rightarrow v = 0.$$

III. Self Check Exercise

1. Let $\dim V = n$, $T : V \rightarrow V$ be a L.T. such that $\text{Range } T = \text{Ker } T$. show that n is even. Prove that $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$, s.t., $T(x_1, x_2) = (x_2, 0)$ is such a L.T.

2. Find range, rank, Ker and nullity of the L.T. defined by

$$(i) T : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ s.t., } T(x_1, x_2) = (x_1 + x_2, x_1) \left[\mathbb{R}^2, 2(0), 0 \right]$$

$$(ii) T : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \text{ s.t., } T(x_1, x_2) = (x_1 + x_2, x_1 - x_2, x_2)$$

$$(iii) T : \mathbb{R}^3 \rightarrow \mathbb{R}^2 \text{ s.t., } T(x_1, x_2, x_3) = (x_1 + x_2, x_1 - x_3)$$

$$\left[(1, 1, 0)(1, -1, 1), 2(0), 0 \right]$$

(iv) The zero and the identity linear transformations

$$(v) T : \mathbb{R}^3 \rightarrow \mathbb{R}^2 \text{ s.t., } T(x_1, x_2, x_3) = (x_1 - x_2, 2x_3 - x_1).$$

3. Find the L.T. From $\mathbb{R}^3 \rightarrow \mathbb{R}^3$ which has its range the subspace spanned by $(1, 0, -1), (1, 2, 2)$.

4. Show that the linear transformation $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ defined by

$$T(x_1, x_2, x_3) = (2x_1, x - x_2, 5x_1 + 4x_2 + x_3) \text{ is invertible.}$$

5. Let T be a L.T. from $\mathbb{R}^7$ onto a 3-dimensional subspace of $\mathbb{R}^5$. Show that $\dim \text{Ker } T = 4$.

LINEAR TRANSFORMATIONS-II

- I. Algebra of Linear Transformations**
- II. Invertible Linear Transformations**
- III. Matrix of a Linear Transformations**
- IV. Self Check Exercise**

I. Algebra of Linear Transformations

I.(a) Sum of Linear Transformations

Let V and W be two vector spaces over the same field F . Let $T : V \rightarrow W$ and $S : V \rightarrow W$ be two linear transformations. We define $T + S$, the sum of T and S by

$$T + S : V \rightarrow W, \text{ s.t.}$$

$$(T + S)v = T(v) + S(v), v \in V$$

Then $T + S$ is also a L.T. from $V \rightarrow W$ as

$$\begin{aligned}(T + S)(\alpha x + \beta y) &= T(\alpha x + \beta y) + S(\alpha x + \beta y) \\ &= \alpha T(x) + \beta T(y) + \alpha S(x) + \beta S(y) \\ &= \alpha(T + S)x + \beta(T + S)y\end{aligned}$$

Again for $\alpha \in F$, we define the product of a L.T. $T : V \rightarrow W$ with α , by $(\alpha T) : V \rightarrow W$ s.t.,

$$(\alpha T)v = \alpha(T(v)).$$

It is easy to see that αT is also a L.T. from $V \rightarrow W$. Let $\text{Hom}(V, W)$ be the set of all linear transformations from $V \rightarrow W$. Then we show $\text{Hom}(V, W)$ forms a vector space over F under the addition and scalar multiplication as defined above.

We have already seen that when $T, S \in \text{Hom}(V, W)$, $\alpha \in F$ then $T + S, \alpha T \in \text{Hom}(V, W)$, thus closure holds for these operations. We verify some of the other conditions in the definition.

$$(T + S)v = T(v) + S(v) = S(v) + T(v) = (S + T)v \text{ for all } v \in V$$

$$\Rightarrow T + S = S + T \text{ for all } S, T \in \text{Hom}(V, W)$$

The map $O : V \rightarrow W$, s.t., $O(v) = 0$ is a L.T. and

$$(T + O)v = T(v) + O(v) = T(v) = (O + T)v \text{ for all } v$$

This O is zero of $\text{Hom}(V, W)$

For any $T \in \text{Hom}(V, W)$, the map $(-T) : V \rightarrow W$, s.t.,

$$(-T)v = -T(v)$$

will be additive inverse of T .

$$\text{Again, } [\alpha(T + S)]v = \alpha[(T + S)v] = \alpha[T(v) + S(v)] = \alpha T(v) + \alpha S(v)$$

$$= (\alpha T)v + (\alpha S)v = (\alpha T + \alpha S)v \text{ for all } v \in V$$

$$\Rightarrow \alpha(T + S) = \alpha T + \alpha S$$

$$[(\alpha\beta)T]v = (\alpha\beta)T(v) = \alpha[\beta T(v)] = [\alpha(\beta T)]v \text{ for all } v$$

$$\Rightarrow (\alpha\beta)T = \alpha(\beta T)$$

$$(1T)v = 1.T(v) = T(v) \text{ for all } v$$

$$\Rightarrow 1 \cdot T = T$$

Hence one notices that $\text{Hom}(V, W)$ forms a vector space over F .

Note : The notation $L(V, W)$ is also used for denoting $\text{Hom}(V, W)$.

I.(b) Product of Linear's Transformations

Definition : Product (composition) of two linear transformations

Let V, W, Z be three vector spaces over a field F

Let $T : V \rightarrow W, S : W \rightarrow Z$ be L.T.

We define $ST : V \rightarrow Z$, s.t.,

$$(ST)v = S(T(v))$$

then ST is a linear transformation (verify!), called product of S and T .

Note : TS may not be defined and even if it is defined it may not equal ST .

II. Linear operator and Linear Functional

Definition : A L.T. $T : V \rightarrow V$ is called a linear operator on V , whereas a L.T. $T : V \rightarrow F$ is called a linear functional. We use notation T^2 for $T.T$ and $T^n = T^{n-1}T$ etc.

Theorem 11 : Let T, T_1, T_2 be linear operators on V , and let $I : V \rightarrow V$ be the identity map $I(v) = v$ for all v (which is clearly a L.T.) then

$$(i) \quad IT = TI = T$$

$$(ii) \quad T(T_1 + T_2) = TT_1 + TT_2$$

$$(T_1 + T_2)T = T_1T + T_2T$$

$$(iii) \quad \alpha(T_1T_2) = (\alpha T_1)T_2 = T_1(\alpha T_2) \quad \alpha \in F$$

$$(iv) \quad T_1(T_2T_3) = (T_1T_2)T_3.$$

Proof : (i) Obvious.

$$\begin{aligned} (ii) \quad [T(T_1 + T_2)]x &= T[(T_1 + T_2)x] = T[T_1(x) + T_2(x)] \\ &= T(T_1(x)) + T(T_2(x)) = TT_1(x) + TT_2(x) \\ &= (TT_1 + TT_2)x \end{aligned}$$

$$\Rightarrow T(T_1 + T_2) = TT_1 + TT_2$$

Other result follows similarly.

$$\begin{aligned} (iii) \quad [\alpha(T_1T_2)]x &= \alpha[(T_1T_2)x] = \alpha[T_1(T_2(x))] \\ &= [(\alpha T_1)T_2]x = (\alpha T_1)[T_2(x)] = \alpha[T_1(T_2(x))] \\ &= [T_1(\alpha T_2)]x = T_1(\alpha T_2)x = T_1(\alpha T_2(x)) = \alpha T_1(T_2(x)) \end{aligned}$$

Hence the result follows.

(iv) Follows easily by definition.

See exercises for the generalised version of above theorem.

Problem 1 : Find the range, Rank, Ker and nullity of the linear transformation

$$T: \mathbb{R}^3 \rightarrow \mathbb{R}^3, \text{ s.t.,}$$

$$T(x, y, z) = (x + z, x + y + 2z, 2x + y + 3z)$$

Solution : Let $(x, y, z) \in \text{Ker } T$ be any element, then

$$T(x, y, z) = (0, 0, 0)$$

$$\Rightarrow (x + z, x + y + 2z, 2x + y + 3z) = (0, 0, 0)$$

$$\Rightarrow x + 0 + z = 0$$

$$x + y + 2z = 0$$

$$2x + y + 3z = 0$$

Giving $x = -z, -z + y + 2z = 0$ i.e., $y = -z$

Thus Ker T consists of all elements of the type $(x, x, -x) = x(1, 1, -1)$ where x is any real no. i.e., Ker T is spanned by $(1, 1, -1)$ which is L.I. Note $(1, 1, -1) \in \text{Ker T}$

Hence $\dim(\text{Ker T}) = 1 = \text{nullity of T}$

Again, from def. of T, we notice elements of the types $(x + z, x + y + 2z, 2x + y + 3z)$ are in Range T.

$$\begin{aligned}\text{Now } (x + z, x + y + 2z, 2x + y + 3z) &= (x + 0 + z, x + y + 2z, 2x + y + 3z) \\ &= (x, x, 2x) + (0, y, y) + (z, 2z, 3z) \\ &= x(1, 1, 2) + y(0, 1, 1) + z(1, 2, 3)\end{aligned}$$

Thus Range T is spanned by $\{(1, 1, 2), (0, 1, 1), (1, 2, 3)\}$

Since $(1, 1, 2) + (0, 1, 1) = (1, 2, 3)$ we find these vectors are L.D. So $\dim \text{Range T} \leq 2$

Again as $(1, 1, 2)$ and $(0, 1, 1)$ are L.I. we find

$$\dim \text{Range T} = 2 = \text{Rank T}.$$

Problem 2 : Find the range, rank, Ker and nullity of the following linear transformations

$$(a) \quad T: \mathbb{R}^2 \rightarrow \mathbb{R}^3 \text{ s.t., } T(x_1, x_2) = (x_1, x_1 + x_2, x_2)$$

$$(b) \quad T: \mathbb{R}^4 \rightarrow \mathbb{R}^3 \text{ s.t., } T(x_1, x_2, x_3, x_4) = (x_1 - x_4, x_2 + x_3, x_3 - x_4)$$

Solution : (a) From definition of T, we notice elements of the type $(x_1, x_1 + x_2, x_2)$ will have pre images in $\mathbb{R}^2$ i.e., elements of this type are in Range T.

$$\begin{aligned}\text{Now } (x_1, x_1 + x_2, x_2) &= (x_1 + 0, x_1 + x_2, 0 + x_2) \\ &= (x_1, x_1, 0) + (0, x_2, x_2) \\ &= x_1(1, 1, 0) + x_2(0, 1, 1)\end{aligned}$$

or that Range T is spanned by $\{(1, 1, 0), (0, 1, 1)\}$ and since

$$\begin{aligned}\alpha_1(1, 1, 0) + \alpha_2(0, 1, 1) &= (0, 0, 0) \\ \Rightarrow \alpha_1 &= \alpha_2 = 0\end{aligned}$$

these are L.I. and thus form a basis of Range T

$$\Rightarrow \text{Rank T} = \dim \text{Range T} = 2.$$

$$\text{Again } (x_1, x_2) \in \text{Ker T} \Rightarrow T(x_1, x_2) = (0, 0, 0)$$

$$\Rightarrow (x_1, x_1 + x_2, x_2) = (0, 0, 0)$$

$$\Rightarrow x_1 = 0, x_1 + x_2 = 0, x_2 = 0$$

$$\Rightarrow x_1 = x_2 = 0$$

$$\Rightarrow \text{Ker } T = \{(0, 0)\}$$

Also then nullity $T = \dim \text{Ker } T = 0$.

(b) From definition of T , we find elements of the type $(x_1 - x_4, x_2 + x_3, x_3 - x_4)$ have pre image in $\mathbb{R}^4$.

Now

$$\begin{aligned} (x_1 - x_4, x_2 + x_3, x_3 - x_4) &= (x_1 + 0 + 0 - x_4, 0 + x_2 + x_3 + 0, 0 + 0 + x_3 - x_4) \\ &= x_1(1, 0, 0) + x_2(0, 1, 0) + x_3(0, 1, 1) + x_4(-1, 0, -1) \end{aligned}$$

or that Range T is spanned by

$$\{(1, 0, 0), (0, 1, 0), (0, 1, 1), (-1, 0, -1)\}$$

Since Range T is a subspace of $\mathbb{R}^3$ which has dim 3 these four elements cannot form basis of Range T .

In fact these are L.D., elements as

$$(-1, 0, -1) + (1, 0, 0) + (0, 1, 0) + (0, 1, 1) = (0, 0, 0)$$

If we consider three members

$$(1, 0, 0), (0, 1, 0), (0, 1, 1)$$

$$\text{we notice } \alpha_1(1, 0, 0) + \alpha_2(0, 1, 0) + \alpha_3(0, 1, 1) = (0, 0, 0)$$

$$\Rightarrow \alpha_i = 0 \text{ for all } i$$

or that $(1, 0, 0), (0, 1, 0), (0, 1, 1)$ are L.I., and hence form basis of Range T

$$\Rightarrow \dim \text{Range } T = 3 = \text{rank of } T$$

one might notice here that as

$$(-1, 0, -1) = -1(1, 0, 0) - 1(0, 1, 0) - 1(0, 1, 1)$$

the elements $(1, 0, 0), (0, 1, 0), (0, 1, 1)$ span Range T

Also then Range $T = \mathbb{R}^3$

$$\text{Again } (x_1, x_2, x_3, x_4) \in \text{Ker } T \Rightarrow T(x_1, x_2, x_3, x_4) = (0, 0, 0)$$

$$\Rightarrow \quad x_1 - x_4 = 0$$

$$x_2 + x_3 = 0$$

$$x_3 - x_4 = 0$$

if we fix x_4 , we get $x_1 = x_4, x_2 = -x_3 = -x_4, x_3 = x_4$

or that elements of the type $(x_4, -x_4, x_4, x_4)$ are in the Ker T

Problem 3 : Let T be a linear operator on V. If $T^2 = 0$, what can you say about the relation of the range of T to the null space of T? Give an example of linear operator T of $\mathbb{R}^2$ such that $T^2 = 0$, but $T \neq 0$.

Solution : $T^2 = 0 \Rightarrow T^2(v) = 0$ for all $v \in V$

$$\Rightarrow T(T(V)) = 0$$

$$\Rightarrow T(v) \in \text{Ker } T \text{ for all } v \in V$$

$$\Rightarrow \text{range } T \subseteq \text{Ker } T.$$

Define $T: \mathbb{R}^2 \rightarrow \mathbb{R}^2$, such that

$$T(x_1, x_2) = (x_2, 0)$$

then T is a linear operator (Verify!)

Since $T(2, 2) = (2, 0) \neq (0, 0)$

$$T \neq 0$$

But $T^2(x_1, x_2) = T(T(x_1, x_2)) = T(x_2, 0) = (0, 0)$

$$\Rightarrow T^2 = 0.$$

Problem 4 : Let T be a linear operator on V and let $\text{Rank } T^2 = \text{Rank } T$ then show that $\text{Rank } T \cap \text{Ker } T = \{0\}$.

Solution : $T: V \rightarrow V, T^2: V \rightarrow V$

$$\text{Rank } T^2 = \dim V - \dim \text{Ker } T^2$$

$$\Rightarrow \dim \text{Ker } T = \dim \text{Ker } T^2$$

We claim $\text{Ker } T = \text{Ker } T^2$

$$x \in \text{Ker } T \Rightarrow T(x) = 0 \Rightarrow T^2(x) = T(0) = 0$$

$$\Rightarrow x \in \text{Ker } T^2 \Rightarrow \text{Ker } T \subseteq \text{Ker } T^2$$

$$\Rightarrow \text{Ker } T = \text{Ker } T^2 \text{ (as they have same dim)}$$

Now

$$x \in \text{Range } T \cap \text{Ker } T \Rightarrow x \in \text{Range } T \text{ and } x \in \text{Ker } T$$

$$\Rightarrow T(x) = 0, x = T(y) \text{ for some } x \in V$$

$$\Rightarrow T(T(y)) = 0$$

$$\Rightarrow T^2(y) = 0$$

$$\Rightarrow y \in \text{Ker } T^2 = \text{Ker } T$$

$$\Rightarrow T(y) = 0 \Rightarrow x = 0$$

$$\Rightarrow \text{Ker } T \cap \text{Range } T = \{0\}.$$

III. Invertible Linear Transformations

We recall that a map $T: V \rightarrow W$ is invertible iff it is 1-1 onto, and inverse of T is the map $T^{-1}: W \rightarrow V$ such that

$$T^{-1}(y) = x \Leftrightarrow T(x) = y$$

We show that inverse of a (1-1 onto) L.T. is also a L.T. Let $T: V \rightarrow W$ be 1-1 onto L.T. and $T^{-1}: W \rightarrow V$ be its inverse.

We have to prove

$$T^{-1}(\alpha w_1 + \beta w_2) = \alpha T^{-1}(w_1) + \beta T^{-1}(w_2) \quad \alpha, \beta \in F, w_1, w_2 \in W$$

Since T is onto, for $w_1, w_2 \in W, \exists v_1, v_2 \in V$ such that $T(v_1) = w_1, T(v_2) = w_2$

$$\Leftrightarrow v_1 = T^{-1}(w_1), v_2 = T^{-1}(w_2)$$

Now

$$T^{-1}(\alpha w_1 + \beta w_2) = T^{-1}(\alpha T(v_1) + \beta T(v_2))$$

$$= T^{-1}(T(\alpha v_1) + T(\beta v_2))$$

$$= T^{-1}(T(\alpha v_1 + \beta v_2))$$

$$= \alpha v_1 + \beta v_2$$

$$= \alpha T^{-1}(w_1) + \beta T^{-1}(w_2).$$

Definition : A L. T. $T: V \rightarrow W$ is called non-singular if $\text{Ker } T = \{0\}$ i.e. if T is 1-1.

Theorem 12 : A linear transformation $T: V \rightarrow W$ is non singular iff T carries each L.I. subset of V onto a L.I. subset of W .

Proof : Let T be non-singular and $\{v_1, v_2, \dots, v_n\}$ be a L.I. subset of V . we show

$\{T(v_1), T(v_2), \dots, T(v_n)\}$ is L.I. subset of W .

$$\text{Now } \alpha_1 T(v_1) + \alpha_2 T(v_2) + \dots + \alpha_n T(v_n) = 0 \quad \alpha_i \in F$$

$$\Rightarrow T(\alpha_1 v_1 + \dots + \alpha_n v_n) = 0$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n \in \text{Ker } T = \{0\}$$

$$\Rightarrow \alpha_1 v_1 + \dots + \alpha_n v_n = 0$$

$$\Rightarrow \alpha_i = 0 \text{ for all } i \text{ as } v_1, v_2, \dots, v_n \text{ are L.I.}$$

Conversely, let $v \in \text{Ker } T$ be any element

$$\text{Then } T(v) = 0$$

$$\Rightarrow \{T(v)\} \text{ is not L.I. in } W$$

$$\Rightarrow v \text{ is not L.I. in } V. \text{ (by hypothesis)}$$

$$\Rightarrow v = 0 \Rightarrow \text{Ker } T = \{0\}$$

$$\Rightarrow T \text{ is non singular.}$$

Theorem 13 : Let $T: V \rightarrow W$ be a L.T. where V and W are two F.D.V.S. with same dimension. Then the following are equivalent.

- (i) T is invertible
- (ii) T is non singular (i.e., T is 1-1)
- (iii) T is onto (i.e. $\text{Range } T = W$)
- (iv) If $\{v_1, v_2, \dots, v_n\}$ is a basis of V then

$$\{T(v_1), T(v_2), \dots, T(v_n)\} \text{ is a basis of } W.$$

Proof : (i) $\Rightarrow$ (ii) follows by definition.

$$(ii) \Rightarrow (iii) \quad T \text{ is non-singular}$$

$$\Rightarrow \text{Ker } T = \{0\}$$

$$\Rightarrow \dim \text{Ker } T = 0$$

Since $\dim \text{Range } T + \dim \text{Ker } T = \dim V$, we get

$$\dim \text{Range } T = \dim V$$

$$\Rightarrow \dim \text{Range } T = \dim W \text{ (given condition)}$$

But $\text{Range } T$ being a subspace of W , we find

$$\text{Range } T = W$$

$$(iii) \Rightarrow (i) \text{ } T \text{ onto means } \text{Range } T = W$$

$$\Rightarrow \dim \text{Range } T = \dim W = \dim V$$

and as $\dim \text{Range } T + \dim \text{Ker } T = \dim V$, we get

$$\dim \text{Ker } T = 0$$

$$\Rightarrow \text{Ker } T = \{0\}$$

or that T is 1-1 and as it is onto T will be invertible.

$$(i) \Rightarrow (iv) \text{ } T \text{ is invertible} \Rightarrow T \text{ is 1-1 onto}$$

i.e., T is an isomorphism.

$$(iv) \Rightarrow (i)$$

Let $\{T(v_1), \dots, T(v_n)\}$ be basis of W where $\{v_1, \dots, v_n\}$ is basis of V . Any $w \in W$ can be put as

$$w = \alpha_1 T(v_1) + \dots + \alpha_n T(v_n)$$

$$= T(\alpha_1 v_1 + \dots + \alpha_n v_n) = T(v) \text{ for some } v \in V$$

$\therefore T$ is onto. Thus (iii) holds.

Hence (i) holds.

Problem 5 : Let T be a linear operator on $\mathbb{R}^3$, defined by

$$T(x_1, x_2, x_3) = (3x_1, x_1 - x_2, 2x_1 + x_2 + x_3)$$

show that T is invertible and find the rule by which T^{-1} is defined.

Solution: $T: \mathbb{R}^3 \rightarrow \mathbb{R}^3$

Let $T(x_1, x_2, x_3) \in \text{Ker } T$ be any element

$$\text{Then } T(x_1, x_2, x_3) = (0, 0, 0)$$

$$\Rightarrow (3x_1, x_1 - x_2, 2x_1 + x_2 + x_3) = (0, 0, 0)$$

$$\Rightarrow 3x_1 = 0, x_1 - x_2 = 0, 2x_1 + x_2 + x_3 = 0$$

$$\Rightarrow x_1 = x_2 = x_3 = 0 \text{ or that } \text{Ker } T = \{(0, 0, 0)\}$$

$\Rightarrow T$ is non singular and thus invertible (See theorem 8)

Now If (z_1, z_2, z_3) be any element of $\mathbb{R}^3$, then (x_1, x_2, x_3) will be its image under T if

$$T(x_1, x_2, x_3) = (z_1, z_2, z_3)$$

$$\Rightarrow 2x_1 = z_1$$

$$x_1 - x_2 = z_2$$

$$2x_1 + x_2 + x_3 = z_3$$

$$\text{Which give } x_1 = \frac{z_1}{3}, x_2 = \frac{z_1}{3} - z_2, z_3 = z_3 - z_1 + z_2$$

Hence $T^{-1} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is defined by

$$T^{-1} : (z_1, z_2, z_3) = \left(\frac{z_1}{3}, \frac{z_1}{3} - z_2, z_3 - z_1 + z_2 \right).$$

Problem 6 : If $T : V \rightarrow V$ is a L.T., such that T is not onto then show that there exists some $0 \neq v$ in V s.t., $T(v) = 0$.

Solution : Since T is not onto, it is not 1-1 (theorem done)

Suppose $\exists$ no $0 \neq v \in V$ s.t. $T(v) = 0$

Then $T(v) = 0$ only when $v = 0$

$\Rightarrow \text{Ker } T = \{0\} \Rightarrow T$ is 1-1, a contradiction.

IV. Matrix of a Linear Transformations

Let $U(F), V(F)$ be vector spaces of dimension n and m respectively, Let

$\beta = \{u_1, \dots, u_n\}, \beta' = \{v_1, \dots, v_m\}$ be their ordered basis respectively. Suppose $T : U \rightarrow V$

is a linear transformation. Since $T(u_1), \dots, T(u_n) \in V$ and $\{v_1, \dots, v_m\}$ spans V , each

$T(u_i)$ is a linear combination of vectors $v_1, \dots, v_m$.

$$\text{Let } T(u_1) = \alpha_{11}v_1 + \dots + \alpha_{m1}v_m$$

$$T(u_2) = \alpha_{12}v_1 + \dots + \alpha_{m2}v_m$$

.....

$$T(u_n) = \alpha_{1n}v_1 + \dots + \alpha_{mn}v_m$$

where each $a_{ij} \in F$. Then the $m \times n$ matrix

$$A = \begin{bmatrix} \alpha_{11} & \alpha_{12} & \dots & \dots & \alpha_{1n} \\ \vdots & \vdots & \dots & \dots & \vdots \\ \vdots & \vdots & \dots & \dots & \vdots \\ \vdots & \vdots & \dots & \dots & \vdots \\ \alpha_{m1} & \alpha_{m2} & \dots & \dots & \alpha_{mn} \end{bmatrix}$$

is called matrix of T with respect to ordered basis β, β' respectively. A is uniquely determined by T as each $\alpha_{ij} \in F$ is uniquely determined. We write

$$A = [T]_{\beta, \beta'}$$

Theorem 14 : $\text{Hom}(U, V) = M_{n \times n}(F)$

Proof : Define $\theta : \text{Hom}(U, V) \rightarrow M_{m \times n}(F)$, s.t.,

$$\theta(T) = [T]_{\beta, \beta'}$$

Where $\beta = \{u_1, \dots, u_n\}, \beta' = \{v_1, \dots, v_m\}$ are ordered basis of U, V respectively. θ is well defined as $[T]_{\beta, \beta'}$ is uniquely determined by T

It is not difficult to verify that θ is a linear transformation.

$$\text{Let } \theta(S) = \theta(T), ST \in H(U, V)$$

$$\text{Then } [S]_{\beta, \beta'} = [T]_{\beta, \beta'}$$

$$\Rightarrow (a_{ij}) = (b_{ij})$$

$$\Rightarrow a_{ij} = b_{ij} \text{ for all } i, j$$

$$\Rightarrow S(u_j) = \sum_{i=1}^m a_{ij}v_i = \sum_{i=1}^m b_{ij}v_i = T(u_j) \text{ for all } j = 1, \dots, n$$

$$\Rightarrow S = T \Rightarrow \theta \text{ is } 1-1$$

Let $A = (a_{ij})_{m \times n} \in M_{m \times n}(F)$. Then $\exists$ a linear transformation $T \in H(U, V)$ s.t.,

$$T(u_j) = \sum_{i=1}^m a_{ij} v_i \quad \text{for } j = 1, \dots, n$$

$$A = [T]_{\beta, \beta'} = \theta(T) \Rightarrow \theta \text{ is onto.}$$

Hence θ is an isomorphism and so $\text{Hom}(U, V) \cong M_{m \times n}(F)$

Cor $\therefore \dim \text{Hom}(U, V) = mn$

Proof : S = set of all $m \times n$ matrices with only one entry 1 and all other entries zero, is a basis of $M_{m \times n}(F)$.

Clearly, $o(S) = mn \Rightarrow \dim M_{m \times n}(F) = mn$

$$\dim \text{Hom}(U, V) = mn.$$

Problem 7 : Let T be a linear operator on C^2 defined by $T(x_1, x_2) = (x_1, 0)$. Let $\beta = \{\epsilon_1 = (1, 0), \epsilon_2 = (0, 1)\}$, $\beta' = \{\alpha_1 = (1, i), \alpha_2 = (-i, 2)\}$ be ordered basis for C^2 . What is the matrix of T relative to the pair β, β' ?

Solution : Now $T(\epsilon_1) = T(1, 0)$
 $= (1, 0)$
 $= a(1, i) + b(-i, 2)$
 $\Rightarrow a - bi = 1$ where $a, b \in C$
 $ai + 2b = 0$
 $\Rightarrow a = 2, b = -i$
 $\Rightarrow T(\epsilon_1) = 2\alpha_1 - i\alpha_2$

Also $T(\epsilon_2) = T(0, 1) = (0, 0) = 0\alpha_1 + 0\alpha_2$

$$\therefore [T]_{\beta\beta'} = \begin{bmatrix} 2 & 0 \\ -i & 0 \end{bmatrix}.$$

Problem 8 : Let T be linear operator on R^3 , the matrix of which in the standard ordered basis is:

$$A = \begin{bmatrix} 1 & 2 & 1 \\ 0 & 1 & 1 \\ -1 & 3 & 4 \end{bmatrix}$$

Find a basis for the range of T and a basis for the null space of T.

Solution : Det $A = 1(4 - 3) - 2(1) + 1(1)$

$$= 1 - 2 + 1 = 0$$

$\therefore$ A is not invertible and so T is not invertible

$$\text{Let } \{\epsilon_1 = (1, 0, 0), \epsilon_2 = (0, 1, 0), \epsilon_3 = (0, 0, 1)\}$$

be standard ordered basis of $\mathbb{R}^3$.

$$\text{Let } (x_1, x_2, x_3) \in \text{Ker } T$$

$$\text{Then } T(x_1, x_2, x_3) = 0$$

$$\Rightarrow T(x_1(1, 0, 0) + x_2(0, 1, 0) + x_3(0, 0, 1)) = 0$$

$$\Rightarrow T(x_1 \epsilon_1 + x_2 \epsilon_2 + x_3 \epsilon_3) = 0$$

$$\Rightarrow x_1 T(\epsilon_1) + x_2 T(\epsilon_2) + x_3 T(\epsilon_3) = 0$$

$$\Rightarrow x_1(1, 0, -1) + x_2(2, 1, 3) + x_3(1, 1, 4) = 0$$

$$\Rightarrow (x_1 + 2x_2 + x_3, x_2 + x_3, -x_1 + 3x_2 + 4x_3) = 0$$

$$\Rightarrow x_1 + 2x_2 + x_3 = 0, x_2 + x_3 = 0, -x_1 + 3x_2 + 4x_3 = 0$$

$$\Rightarrow x_1 + x_2 = 0, x_2 + x_3 = 0$$

$$\Rightarrow (x_1, x_2, x_3) = (-x_2, x_2, -x_3)$$

$$= x_2(-1, 1, -1)$$

$$\Rightarrow \text{every element in Ker } T \text{ is multiple of } (-1, 1, -1)$$

$$\Rightarrow \text{Ker } T \text{ is spanned by } (-1, 1, -1)$$

Since $(-1, 1, -1) \neq 0$, $\{(-1, 1, -1)\}$ is basis of Ker T.

$$\therefore \dim \text{Ker } T = 1 \Rightarrow \dim \text{Range } T = 2$$

$$\text{Since } T \epsilon_1 = (1, 0, -1)$$

$$T \epsilon_2 = (2, 1, 3)$$

belong to Range T and $aT \in_1 + bT \in_2 = 0$

we find $a(1, 0, -1) + b(2, 1, 3) = 0$

$$\Rightarrow b = 0, a = 0$$

$\Rightarrow \{T \in_1, T \in_2\}$ is a linearly independent set in Range T. As $\dim \text{Range T} = 2$,

$\{(1, 0, -1), (2, 1, 3)\}$ is a basis of Range T.

Problem 9 : Let T be a linear operator on F^n and let A be the matrix of T in the standard ordered basis for F^n . Let W be the subspace of F^n spanned by the column vectors of A. Find a relation between W and T.

Solution : $T: F^n \rightarrow F^n$

Let $\beta = \{e_1 = (1, 0, 0, \dots, 0), \dots, e_n = (0, 0, \dots, 1)\}$ be the standard ordered basis of F^n and let

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & \dots & \vdots \\ a_{n1} & a_{n2} & \dots & a_{nn} \end{bmatrix}$$

$$\text{thus } T(e_1) = a_{11}e_1 + a_{21}e_2 + \dots + a_{n1}e_n$$

$$T(e_2) = a_{12}e_1 + a_{22}e_2 + \dots + a_{n2}e_n$$

$$\dots \quad \dots \quad \dots$$

$$T(e_n) = a_{1n}e_1 + a_{2n}e_2 + \dots + a_{nn}e_n$$

and also W is spanned by

$$\{(a_{11}, a_{21}, \dots, a_{n1}), (a_{12}, a_{22}, \dots, a_{n2}), \dots, (a_{1n}, a_{2n}, \dots, a_{nn})\}$$

We claim $T: F^n \rightarrow W$ is onto L.T.

For any $x \in F^n$, $x = \alpha_1 e_1 + \alpha_2 e_2 + \dots + \alpha_n e_n$

$$\Rightarrow T(x) = \alpha_1 T(e_1) + \alpha_2 T(e_2) + \dots + \alpha_n T(e_n)$$

$$\Rightarrow T(x) \in W \text{ as } T(e_1), T(e_2), \dots, T(e_n) \in W$$

Again, for any $w \in W$, $w = \beta_1 T(e_1) + \beta_2 T(e_2) + \dots + \beta_n T(e_n)$

$$= T(\beta_1 e_1 + \beta_2 e_2 + \dots + \beta_n e_n)$$

showing that T is onto

$$\Rightarrow \text{Range } T = W \Rightarrow \dim \text{Range } T = \dim W$$

or that $\text{rank of } T = \dim W$

which is the required relation between T and W .

V. Self Check Exercise

1. Let T be the linear transformation from $\mathbb{R}^3$ into $\mathbb{R}^2$ defined by

$$T(x_1, x_2, x_3) = (x_1 + x_2, 2x_1 - x_3)$$

(i) If β, β' are standard ordered basis for $\mathbb{R}^3$ and $\mathbb{R}^2$ respectively, find $[T]_{\beta, \beta'}$.

$$(ii) \text{ If } \beta = \{\alpha_1 = (1, 0, -1), \alpha_2 = (1, 1, 1), \alpha_3 = (1, 0, 0)\}$$

$$\beta' = \{\beta_1 = (0, 1), \beta_2 = (1, 0)\}.$$

2. Let T be the linear operator on $\mathbb{R}^3$, the matrix of which in the standard

ordered basis is $A = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \\ 0 & 1 & 1 \end{bmatrix}$

Find a basis of $\text{Range } T$ and $\text{Ker } T$.

3. Show that $T: \mathbb{R}^3 \rightarrow \mathbb{R}^2$ s.t, $T(x, t, z) = (y, x)$ is a L.T. Find the matrix representation of T for the standard ordered basis for $\mathbb{R}^3$ and $\{(0, 1), (2, 3)\}$ of $\mathbb{R}^2$.